

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 April 2002 (25.04.2002)

PCT

(10) International Publication Number
WO 02/32893 A2

(51) International Patent Classification⁷: C07D 401/00

Mwangi, Wa; 45 Snyder Road, Fords, NJ 08863 (US). **PI-
WINSKI, John, J.**; 6 Saddle Ridge Drive, Clinton Town-
ship, NJ 08833 (US).

(21) International Application Number: PCT/US01/32151

(22) International Filing Date: 15 October 2001 (15.10.2001)

(74) Agent: **JEANETTE, Henry, C.**; Schering-Plough Corpo-
ration, Patent Department - K-6-1 1990, 2000 Galloping
Hill Road, Kenilworth, NJ 07033-0530 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/240,901 17 October 2000 (17.10.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ,
DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU,
ID, IL, IN, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LU,
LV, MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PH, PL,
PT, RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,
UZ, VN, YU, ZA.

(71) Applicant: **SCHERING CORPORATION** [US/US];
Patent Department - K-6-1 1990, 2000 Galloping Hill
Road, Kenilworth, NJ 07033-0530 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

(72) Inventors: **ASLANIAN, Robert, G.**; 144 Philip Drive,
Rockaway, NJ 07866 (US). **SHIH, Neng-Yang**; 1 Maple
Drive, North Caldwell, NJ 07006 (US). **TING, Pauline,**
C.; 74 Delwick Lane, New Providence, NJ 07974 (US).
BERLIN, Michael, Y.; 14 Hendrick Road, Flemington,
NJ 08822 (US). **ROSENBLUM, Stuart, B.**; 16 Steven
Terrace, West Orange, NJ 07052 (US). **MC CORMICK,**
Kevin, D.; 5 Pace Drive, Edison, NJ 08820 (US). **TOM,**
Wing, C.; 133 Cedar Grove Parkway, Cedar Grove, NJ
07009 (US). **BOYCE, Christopher, W.**; 71 Bonnell Street,
Flemington, NJ 08822 (US). **MANGIARACINA, Pietro;**
4 Montclair Avenue, Monsey, NY 10952 (US). **MUTAH,**


Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



(54) Title: NOVEL NON-IMIDAZOLE COMPOUNDS



(57) Abstract: Disclosed are novel compounds of the formula: (I) Also disclosed are pharmaceutical compositions comprising the compounds of Formula (I). Also disclosed are methods of treating various diseases or conditions, such as, for example, allergy, allergy-induced airway responses, and congestion (e.g., nasal congestion) using the compounds of Formula (I). Also disclosed are methods of treating various diseases or conditions, such as,

for example, allergy, allergy-induced airway responses, and congestion (e.g., nasal congestion) using the compounds of Formula (I) in combination with a H1 receptor antagonist.

5

NOVEL NON-IMIDAZOLE COMPOUNDS

10 REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application Serial No. 60/240901 filed October 17, 2000.

BACKGROUND OF THE INVENTION

15 WO 95/14007 published May 26, 1995 discloses H₃ receptor antagonists of the imidazole type.

WO99/24405 published May 20, 1999 discloses H₃ receptor ligands of the imidazole type.


20 US 5,869,479 issued February 9, 1999 discloses compositions for the treatment of the symptoms of allergic rhinitis using a combination of at least one histamine H₁ receptor antagonist and at least one histamine H₃ receptor antagonist.

In view of the art's interest in compounds which affect H₃ receptors, novel compounds that are antagonists of H₃ receptors would be a welcome contribution to the art. This invention provides just such a contribution.

25

Summary of the Invention

The present invention provides novel compounds of structure I.



30 or a pharmaceutically acceptable salt or solvate thereof, wherein:

(1) R¹ is selected from:


- 2 -

- (a) aryl;
- (b) heteroaryl;
- (c) heterocycloalkyl
- (d) alkyl;
- (e) cycloalkyl; or
- (f) alkylaryl;

wherein said R^1 groups are optionally substituted with 1 to 4 substituents independently selected from:

- (1) halogen (e.g., Br, F, or Cl, preferably F or Cl);
- (2) hydroxyl (i.e., -OH);
- (3) lower alkoxy (e.g., C_1 to C_6 alkoxy, preferably C_1 to C_4 alkoxy, most preferably C_1 to C_2 alkoxy, more preferably methoxy);
- (4) $-CF_3$;
- (5) CF_3O- ;
- (6) $-NR^4R^5$;
- (7) phenyl;
- (8) $-NO_2$;
- (9) $-CO_2R^4$;
- (10) $-CON(R^4)_2$ wherein each R^4 is the same or different;
- (11) $-S(O)_mN(R^{20})_2$ wherein each R^{20} is the same or different H or alkyl group, preferably C_1 to C_4 alkyl, most preferably C_1 - C_2 alkyl, and more preferably methyl;
- (12) $-CN$; or
- (13) alkyl; or

- (2) R^1 and X taken together form a group selected from:




- (3) X is selected from: $=C(O)$, $=C(NOR^3)$, $=C(NNR^4R^5)$,



- 3 -

- (4) M^1 is carbon;
- (5) M^2 is selected from C or N;
- (6) M^3 and M^4 are independently selected from C or N;
- (7) Y is selected from: is $-\text{CH}_2-$, $=\text{C}(\text{O})$, $=\text{C}(\text{NOR}^{20})$ (wherein R^{20} is as
 5 defined above), or $=\text{C}(\text{S})$;
- (8) Z is a $\text{C}_1 - \text{C}_6$ alkyl group;
- (9) R^2 is a five or six-membered heteroaryl ring, said six-membered heteroaryl ring comprising 1 or 2 nitrogen atoms with the remaining ring atoms being carbon, and said five-membered heteroaryl ring containing 1 or 2 heteroatoms
 10 selected from: nitrogen, oxygen, or sulfur with the remaining ring atoms being carbon; said five or six membered heteroaryl rings being optionally substituted with 1 to 3 substituents independently selected from: halogen, hydroxyl, lower alkyl, lower alkoxy, $-\text{CF}_3$, $\text{CF}_3\text{O}-$, $-\text{NR}^4\text{R}^5$, phenyl, $-\text{NO}_2$, $-\text{CO}_2\text{R}^4$, $-\text{CON}(\text{R}^4)_2$ wherein each R^4 is the same or different, $-\text{CH}_2\text{NR}^4\text{R}^5$, $-(\text{N})\text{C}(\text{NR}^4\text{R}^5)_2$, or $-\text{CN}$;
- (10) R^3 is selected from:
- (a) hydrogen;
- (b) $\text{C}_1 - \text{C}_6$ alkyl;
- (c) aryl;
- (d) heteroaryl;
- (e) heterocycloalkyl;
- (f) arylalkyl (e.g., $\text{aryl}(\text{C}_1 \text{ to } \text{C}_4)\text{alkyl}$, e.g., $-(\text{CH}_2)_w\text{aryl}$ wherein w is 1
 20 to 4, preferably 1 or 2, and most preferably 1, such as, for example $-\text{CH}_2\text{phenyl}$ or $-\text{CH}_2\text{substituted phenyl}$);
- (g) $-(\text{CH}_2)_e-\text{C}(\text{O})\text{N}(\text{R}^4)_2$ wherein each R^4 is the same or different,
- (h) $-(\text{CH}_2)_e-\text{C}(\text{O})\text{OR}^4$;
- (i) $-(\text{CH}_2)_e-\text{C}(\text{O})\text{R}^{30}$ wherein R^{30} is a heterocycloalkyl group, such as,
 25 for example, morpholinyl, piperidinyl, piperazinyl or pyrrolidinyl, including



- (j) $-\text{CF}_3$; or
- (k) $-\text{CH}_2\text{CF}_3$;
- 30

- 4 -

wherein said aryl, heteroaryl, heterocycloalkyl, and the aryl portion of said arylalkyl are optionally substituted with 1 to 3 (preferably 1) substituents selected from:

halogen (e.g., F or Cl), -OH, -OCF₃, -CF₃, -CN, -N(R⁴⁵)₂, -CO₂R⁴⁵, or -C(O)N(R⁴⁵)₂,

wherein each R⁴⁵ is independently selected from: H, alkyl, alkylaryl, or alkylaryl

5 wherein said aryl moiety is substituted with 1 to 3 substituents independently selected from -CF₃, -OH, halogen, alkyl, -NO₂, or -CN;

(11) R⁴ is selected from: hydrogen, C₁ - C₆ alkyl, aryl, alkylaryl, said aryl and alkylaryl groups being optionally substituted with 1 to 3 substituents selected from:

halogen, -CF₃, -OCF₃, -OH, -N(R⁴⁵)₂, -CO₂R⁴⁵, -C(O)N(R⁴⁵)₂, or -CN; wherein R⁴⁵ is

10 as defined above;

(12) R⁵ is selected from: hydrogen, C₁ - C₆ alkyl, -C(O)R⁴, -C(O)₂R⁴, or -C(O)N(R⁴)₂ wherein each R⁴ is independently selected, and R⁴ is as defined above;

(13) or R⁴ and R⁵ taken together with the nitrogen atom to which they are bound forms a five or six membered heterocycloalkyl ring (e.g., morpholine);

15 (14) R⁶ is selected from: alkyl, aryl, alkylaryl, halogen, hydroxyl, lower alkoxy, -CF₃, CF₃O-, -NR⁴R⁵, phenyl, -NO₂, -CO₂R⁴, -CON(R⁴)₂ wherein each R⁴ is the same or different, or -CN;

(15) R¹² is selected from: alkyl, hydroxyl, alkoxy, or fluoro;

(16) R¹³ is selected from: alkyl, hydroxyl, alkoxy, or fluoro;

20 (17) a (subscript for R¹²) is 0 to 2;

(18) b (subscript for R¹³) is 0 to 2;

(19) c (subscript for R⁶) is 0 to 2;

(20) e is 0 to 5;

(21) m is 1 or 2;

25 (22) n is 1, 2 or 3; and

(23) p is 1, 2 or 3, with the proviso that when M³ and M⁴ are both nitrogen, then p is 2 or 3 (i.e., p is not 1 when M³ and M² are both nitrogen).

30 This invention also provides a pharmaceutical composition comprising an effective amount of compound of Formula I, and a pharmaceutically acceptable carrier.

This invention further provides a method of treating: allergy, allergy-induced airway (e.g., upper airway) responses, congestion (e.g., nasal congestion),

hypotension, cardiovascular disease, diseases of the GI tract, hyper and hypo motility and acidic secretion of the gastro-intestinal tract, obesity, sleeping disorders (e.g., hypersomnia, somnolence, and narcolepsy), disturbances of the central nervous system, attention deficit hyperactivity disorder ADHD), hypo and hyperactivity of the central nervous system (for example, agitation and depression), and other CNS disorders (such as Alzheimer's, schizophrenia, and migraine) comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a compound of Formula I.

This invention further provides a method of treating: allergy comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a compound of Formula I.

This invention further provides a method of treating: allergy-induced airway (e.g., upper airway) responses comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a compound of Formula I.

This invention further provides a method of treating: congestion (e.g., nasal congestion) comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a compound of Formula I.

This invention further provides a pharmaceutical composition comprising an effective amount of a compound of Formula I, and an effective amount of a H_1 receptor antagonist in combination with a pharmaceutically acceptable carrier.

This invention further provides a method of treating: allergy, allergy-induced airway (e.g., upper airway) responses, and congestion (e.g., nasal congestion) comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a compound of Formula I in combination with an effective amount of an H_1 receptor antagonist.

This invention further provides a method of treating: allergy comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a compound of Formula I in combination with an effective amount of an H_1 receptor antagonist.

This invention further provides a method of treating: allergy-induced airway (e.g., upper airway) responses comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a

- 6 -

compound of Formula I in combination with an effective amount of an H₁ receptor antagonist.

This invention further provides a method of treating: congestion (e.g., nasal congestion) comprising administering to a patient in need of such treatment (e.g., a mammal, such as a human being) an effective amount of a compound of Formula I in combination with an effective amount of an H₁ receptor antagonist.

DETAILED DESCRIPTION OF THE INVENTION

As used herein, the following terms have the following meanings, unless indicated otherwise:

alkyl-(including the alkyl portions of alkoxy and alkylaryl)-represents straight and branched carbon chains and contains from one to twenty carbon atoms, preferably one to six carbon atoms;

alkylaryl-represents an alkyl group, as defined above, bound to an aryl group, as defined below, wherein said aryl group is bound to the rest of the molecule;

aryl (including the aryl portion of alkylaryl)-represents a carbocyclic group containing from 6 to 15 carbon atoms and having at least one aromatic ring (e.g., aryl is a phenyl ring), with all available substitutable carbon atoms of the carbocyclic group being intended as possible points of attachment;

arylalkyl-represents an aryl group, as defined above, bound to an alkyl group, as defined above, wherein said alkyl group is bound to the rest of the molecule;


cycloalkyl-represents saturated carbocyclic rings of from 3 to 20 carbon atoms, preferably 3 to 7 carbon atoms;

halo (halogen)-represents fluoro, chloro, bromo and iodo;

heteroaryl-represents cyclic groups, having at least one heteroatom selected from O, S or N, said heteroatom interrupting a carbocyclic ring structure and having a sufficient number of delocalized pi electrons to provide aromatic character, with the aromatic heterocyclic groups preferably containing from 2 to 14 carbon atoms; examples include but are not limited to isothiazolyl, isoxazolyl, furazanyl, triazolyl, thiazolyl, thienyl, furanyl (furyl), pyrrolyl, pyrazolyl, pyranyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyridyl (e.g., 2-, 3-, or 4-pyridyl), pyridyl N-oxide (e.g., 2-, 3-, or 4-pyridyl N-oxide), triazinyl, pteridinyl, indolyl (benzopyrrolyl), pyridopyrazinyl, isoquinolyl,

- 7 -

quinolinyl, quinoxolinyl, naphthyridinyl, wherein said pyridyl N-oxide can be represented as:



heterocycloalkyl-represents a saturated, carbocyclic ring containing from 3 to 15 carbon atoms, preferably from 4 to 6 carbon atoms, which carbocyclic ring is interrupted by 1 to 3 hetero groups selected from -O-, -S- or -NR⁴⁰- wherein R⁴⁰ represents C₁ to C₆ alkyl, arylalkyl, -C(O)R⁴, -C(O)OR⁴, or -C(O)N(R⁴⁵)₂ (wherein R⁴⁵ is as defined above, and each R⁴⁵ is independently selected); examples include but are not limited to 2- or 3-tetrahydrofuranyl, 2- or 3- tetrahydrothienyl, 2-, 3- or 4-piperidinyl, 2- or 3-pyrrolidinyl, 2- or 3-piperiziny, 2- or 4-dioxanyl, 1,3-dioxolanyl, 1,3,5-trithianyl, pentamethylene sulfide, perhydroisoquinolinyl, decahydroquinolinyl, trimethylene oxide, azetidiny, 1-azacycloheptanyl, 1,3-dithianyl, 1,3,5-trioxanyl, morpholinyl, thiomorpholinyl, 1,4-thioxanyl, and 1,3,5-hexahydrotriazinyl, thiazolidinyl, tetrahydropyranyl;


lower alkyl-represents an alkyl group, as defined above, that comprises 1 to 6 carbon atoms, preferably 1-4 carbon atoms;

lower alkoxy-represents an alkoxy group whose alkyl moiety comprises 1 to 6 carbon atoms, preferably 1-4 carbon atoms;

=C(O)-represents




=C(NOR³)-represents




wherein (1) represents a mixture of oxime isomers; (2) represents one geometric isomer of the oxime wherein the -OR³ group is on the same side of the double bond as the group to the left of the carbon atom; (3) represents one geometric isomer of the

- 8 -


oxime wherein the $-OR^3$ group is on the same side of the double bond as the group to the right of the carbon atom; and (1) can also be represented as:




$=C(NNR^4R^5)$ represents



and represents a mixture of the isomers




$-(N)C(NR^4R^5)_2$ represents




10

(N) in the structure



represents a nitrogen atom that is located at one of the 4 non-fused positions of the ring, i.e., positions 1, 2, 3 or 4 indicated below:



15

AcOH-represents acetic acid;

t-BOC-represents t-butyloxycarbonyl;

Ci/mmol-represents curie/mmol (a measure of specific activity);

m-CPBA-represents m-chloroperbenzoic acid;

CSA-represents camphorsulfonic acid;

20

CBZ-represents carbonylbenzyloxy $(-C(O)OCH_2C_6H_5)$;

- 9 -

- DBU-represents 1,8-diazabicyclo[5.4.0]undec-7-ene;
DBN-represents 1,5-diazabicyclo[4.3.0]non-5-ene;
DCC-represents dicyclohexylcarbodiimide;
Dibal-H-represents diisobutylaluminum hydride;
5 DIPEA-represents N,N-diisopropylethylamine;
DMAP-represents 4-(dimethylamino)pyridine;
DEC-represents 2-diethylaminoethyl chloride hydrochloride;
DMF-represents dimethylformamide;
EDCI-represents 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide;
10 EtOAc-represents ethyl acetate;
EtOH-represents ethanol;
Fmoc-represents 9-fluorenylmethoxycarbonyl;
HOBT-represents 1-hydroxybenzotriazole;
HPLC-represents high performance liquid chromatography;
15 HRMS-represents high resolution mass spectrometry;
K_i-represents inhibition constant for substrate/receptor complex;
LAH-lithium aluminum hydride;
LDA-represents lithium diisopropylamide;
LRMS-represents low resolution mass spectrometry;
20 MeOH-represents methanol;
NaBH(OAc)₃-represents sodium triacetoxymborohydride;
NaBH₄-represents sodium borohydride;
NaBH₃CN-represents sodium cyanoborohydride;
NaHMDS-represents sodium hexamethyl disilylazide;
25 nM-represents nanomolar;
pA₂-represents -logEC₅₀, as defined by J. Hey, Eur. J. Pharmacol., (1995),
Vol. 294, 329-335;
PCC-represents pyridinium chlorochromate ;
PyBOP-represents benzotriazole-1-yl-oxy-trispyrrolidino-phosphonium
30 hexafluorophosphate;
TEMPO-represents 2,2,6,6-tetramethyl-1-piperidinyloxy, free radical ;
TFA-represents trifluoroacetic acid;
TMAD-represents N,N,N',N'-tetramethylazodicarboxamide;

- 10 -

TMEDA-represents tetramethylethylenediamine;

Tr-represents triphenylmethyl;

Tris-represents tris(hydroxymethyl)aminomethane;and

p-TsOH-represents p-toluenesulfonic acid.

5

Also, as used herein, "upper airway" usually means the upper respiratory system—i.e., the nose, throat, and associated structures.

Also, as used herein, "effective amount" generally means a therapeutically effective amount.

10

Lines drawn into the rings indicate that the indicated bond may be attached to any of the substitutable ring carbon atoms.

Certain compounds of the invention may exist in different isomeric (e.g., enantiomers, diastereoisomers and geometric) forms. The invention contemplates all such isomers both in pure form and in admixture, including racemic mixtures. Enol forms are also included.

15

The compounds of this invention are ligands for the histamine H₃ receptor. The compounds of this invention can also be described as antagonists of the H₃ receptor, or as H₃ antagonists.

The compounds of the invention are basic and form pharmaceutically acceptable salts with organic and inorganic acids. Examples of suitable acids for such salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, oxalic, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those skilled in the art. The salts are prepared by contacting the free base form with a sufficient amount of the desired acid to produce a salt in the conventional manner. The free base forms may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous sodium hydroxide, potassium carbonate, ammonia and sodium bicarbonate. The free base forms differ from their corresponding salt forms somewhat in certain physical properties, such as solubility in polar solvents, but the salts are otherwise equivalent to their corresponding free base forms for purposes of this invention.

20

25

30

The compounds of Formula I can exist in unsolvated as well as solvated forms, including hydrated forms, e.g., hemi-hydrate. In general, the solvated forms, with

- 11 -

pharmaceutically acceptable solvents such as water, ethanol and the like are equivalent to the unsolvated forms for purposes of the invention.

The compounds of this invention can be combined with an H₁ receptor antagonist (i.e., the compounds of this invention can be combined with an H₁ receptor antagonist in a pharmaceutical composition, or the compounds of this invention can be administered with H₁ receptor antagonist).

Numerous chemical substances are known to have histamine H₁ receptor antagonist activity. Many useful compounds can be classified as ethanolamines, ethylenediamines, alkylamines, phenothiazines or piperidines. Representative H₁ receptor antagonists include, without limitation: astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine (also known as SCH-34117), diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast, pyrilamine, promethazine, terfenadine, tripeleminamine, temelastine, trimeprazine and triprolidine. Other compounds can readily be evaluated to determine activity at H₁ receptors by known methods, including specific blockade of the contractile response to histamine of isolated guinea pig ileum. See for example, WO98/06394 published February 19, 1998.

Thus, in the methods of this invention wherein a compound of Formula I is combined with an effective amount of an H₁ receptor antagonist, said H₁ receptor antagonist is selected from: astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast, pyrilamine, promethazine, terfenadine, tripeleminamine, temelastine, trimeprazine or triprolidine.

Also, in the methods of this invention wherein a compound of Formula I is combined with an effective amount of an H₁ receptor antagonist, said H₁ receptor antagonist is selected from: astemizole, azatadine, azelastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, carebastine, descarboethoxyloratadine,

- 12 -

diphenhydramine, doxylamine, ebastine, fexofenadine, loratadine, levocabastine, mizolastine, norastemizole, or terfenadine.

Also, in the methods of this invention wherein a compound of Formula I is combined with an effective amount of an H₁ receptor antagonist, said H₁ receptor antagonist is selected from: azatadine, brompheniramine, cetirizine,
5 chlorpheniramine, carebastine, descarboethoxyloratadine (also known as SCH-34117), diphenhydramine, ebastine, fexofenadine, loratadine, or norastemizole.

Also, in the methods of this invention wherein a compound of Formula I is combined with an effective amount of an H₁ receptor antagonist, said H₁ receptor antagonist is loratadine.
10

Also, in the methods of this invention wherein a compound of Formula I is combined with an effective amount of an H₁ receptor antagonist, said H₁ receptor antagonist is descarboethoxyloratadine.

Also, in the methods of this invention wherein a compound of Formula I is combined with an effective amount of an H₁ receptor antagonist, said H₁ receptor antagonist is fexofenadine.
15

Also, in the methods of this invention wherein a compound of Formula I is combined with an effective amount of an H₁ receptor antagonist, said H₁ receptor antagonist is cetirizine.

Preferably, in the above methods, allergy-induced airway responses are treated.
20

Also, preferably, in the above methods, allergy is treated.

Also, preferably, in the above methods, nasal congestion is treated.

Preferably, in the above methods using a combination of a compound of Formula I (H₃ antagonist) and an H₁ antagonist, the H₁ antagonist is selected from: loratadine, descarboethoxyloratadine, fexofenadine or cetirizine. Most preferably the H₁ antagonist is loratadine or descarboethoxyloratadine.
25

In the methods of this invention wherein a combination of an H₃ antagonist of this invention (compound of Formula I) is administered with a H₁ antagonist, the antagonists can be administered simultaneously, consecutively (one after the other within a relatively short period of time), or sequentially (first one and then the other over a period of time). In general, when the antagonists are administered
30

- 13 -

consecutively or sequentially, the H₃ antagonist of this invention (compound of Formula I) is administered first.

Thus, one embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 32 and a pharmaceutically acceptable carrier.

Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 54 and a pharmaceutically acceptable carrier.

Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 55 and a pharmaceutically acceptable carrier.

Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 253A and a pharmaceutically acceptable carrier.

Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 287 and a pharmaceutically acceptable carrier.

Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 320 and a pharmaceutically acceptable carrier.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, congestion, hypotension, cardiovascular disease, hypotension, diseases of the GI tract, hyper and hypo motility and acidic secretion of the gastro-intestinal tract, obesity, sleeping disorders, disturbances of the central nervous system, attention deficit hyperactivity disorder, hypo and hyperactivity of the central nervous system, Alzheimer's disease, schizophrenia, and migraine comprising administering to a patient in need of such treatment an effective amount of Compound 32.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, congestion, hypotension, cardiovascular disease, hypotension, diseases of the GI tract, hyper and hypo motility and acidic secretion of the gastro-intestinal tract, obesity, sleeping disorders, disturbances of the central nervous system, attention deficit hyperactivity disorder, hypo and hyperactivity

- 14 -

of the central nervous system, Alzheimer's disease, schizophrenia, and migraine comprising administering to a patient in need of such treatment an effective amount of Compound 54.

Another embodiment of this invention is directed to a method of treating:
5 allergy, allergy-induced airway responses, congestion, hypotension, cardiovascular disease, hypotension, diseases of the GI tract, hyper and hypo motility and acidic secretion of the gastro-intestinal tract, obesity, sleeping disorders, disturbances of the central nervous system, attention deficit hyperactivity disorder, hypo and hyperactivity of the central nervous system, Alzheimer's disease, schizophrenia, and migraine
10 comprising administering to a patient in need of such treatment an effective amount of Compound 55.

Another embodiment of this invention is directed to a method of treating:
allergy, allergy-induced airway responses, congestion, hypotension, cardiovascular disease, hypotension, diseases of the GI tract, hyper and hypo motility and acidic
15 secretion of the gastro-intestinal tract, obesity, sleeping disorders, disturbances of the central nervous system, attention deficit hyperactivity disorder, hypo and hyperactivity of the central nervous system, Alzheimer's disease, schizophrenia, and migraine comprising administering to a patient in need of such treatment an effective amount of Compound 253A.

20 Another embodiment of this invention is directed to a method of treating:
allergy, allergy-induced airway responses, congestion, hypotension, cardiovascular disease, hypotension, diseases of the GI tract, hyper and hypo motility and acidic secretion of the gastro-intestinal tract, obesity, sleeping disorders, disturbances of the central nervous system, attention deficit hyperactivity disorder, hypo and hyperactivity
25 of the central nervous system, Alzheimer's disease, schizophrenia, and migraine comprising administering to a patient in need of such treatment an effective amount of Compound 287.

Another embodiment of this invention is directed to a method of treating:
allergy, allergy-induced airway responses, congestion, hypotension, cardiovascular
30 disease, hypotension, diseases of the GI tract, hyper and hypo motility and acidic secretion of the gastro-intestinal tract, obesity, sleeping disorders, disturbances of the central nervous system, attention deficit hyperactivity disorder, hypo and hyperactivity of the central nervous system, Alzheimer's disease, schizophrenia, and migraine

- 15 -

comprising administering to a patient in need of such treatment an effective amount of Compound 320.

Another embodiment of this invention is directed to a method of treating allergy-induced airway responses comprising administering to a patient in need of
5 such treatment an effective amount of Compound 32.

Another embodiment of this invention is directed to a method of treating allergy-induced airway responses comprising administering to a patient in need of such treatment an effective amount of Compound 54.

Another embodiment of this invention is directed to a method of treating
10 allergy-induced airway responses comprising administering to a patient in need of such treatment an effective amount of Compound 55.

Another embodiment of this invention is directed to a method of treating allergy-induced airway responses comprising administering to a patient in need of such treatment an effective amount of Compound 253A.

15 Another embodiment of this invention is directed to a method of treating allergy-induced airway responses comprising administering to a patient in need of such treatment an effective amount of Compound 287.

Another embodiment of this invention is directed to a method of treating allergy-induced airway responses comprising administering to a patient in need of
20 such treatment an effective amount of Compound 320.

Another embodiment of this invention is directed to a method of treating allergy or nasal congestion comprising administering to a patient in need of such treatment an effective amount of Compound 32.

Another embodiment of this invention is directed to a method of treating allergy
25 or nasal congestion comprising administering to a patient in need of such treatment an effective amount of Compound 54.

Another embodiment of this invention is directed to a method of treating allergy or nasal congestion comprising administering to a patient in need of such treatment an effective amount of Compound 55.

30 Another embodiment of this invention is directed to a method of treating allergy or nasal congestion comprising administering to a patient in need of such treatment an effective amount of Compound 253A.

- 16 -

Another embodiment of this invention is directed to a method of treating allergy or nasal congestion comprising administering to a patient in need of such treatment an effective amount of Compound 287.

Another embodiment of this invention is directed to a method of treating allergy
5 or nasal congestion comprising administering to a patient in need of such treatment an effective amount of Compound 320.

Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 32, and an effective amount of H₁ receptor antagonist, and a pharmaceutically effective carrier.

10 Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 54, and an effective amount of H₁ receptor antagonist, and a pharmaceutically effective carrier. Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 55, and an effective amount of H₁ receptor
15 antagonist, and a pharmaceutically effective carrier.

Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 253A, and an effective amount of H₁ receptor antagonist, and a pharmaceutically effective carrier.

Another embodiment of this invention is directed to a pharmaceutical
20 composition comprising an effective amount of Compound 287, and an effective amount of H₁ receptor antagonist, and a pharmaceutically effective carrier. Another embodiment of this invention is directed to a pharmaceutical composition comprising an effective amount of Compound 320, and an effective amount of H₁ receptor antagonist, and a pharmaceutically effective carrier.

25 Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 32 in combination with an effective amount of an H₁ receptor antagonist.

Another embodiment of this invention is directed to a method of treating:
30 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 54 in combination with an effective amount of an H₁ receptor antagonist.

- 17 -

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of a Compound 55 in combination with an effective amount of an H₁ receptor antagonist. Another
5 embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 253A in combination with an effective amount of an H₁ receptor antagonist.

Another embodiment of this invention is directed to a method of treating:
10 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 287 in combination with an effective amount of an H₁ receptor antagonist.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering
15 to a patient in need of such treatment an effective amount of Compound 320 in combination with an effective amount of an H₁ receptor antagonist.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 32 in
20 combination with an effective amount of an H₁ receptor antagonist selected from: astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine,
25 meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast, pyrilamine, promethazine, terfenadine, tripeleminamine, temelastine, trimeprazine or triprolidine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering
30 to a patient in need of such treatment an effective amount of Compound 54 in combination with an effective amount of an H₁ receptor antagonist selected from: astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine,

- 18 -

descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast, pyrilamine, promethazine, terfenadine, tripelennamine, temelastine, trimeprazine or
5 triprolidine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 55 in combination with an effective amount of an H₁ receptor antagonist selected from:
10 astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast,
15 pyrilamine, promethazine, terfenadine, tripelennamine, temelastine, trimeprazine or triprolidine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 253A in combination with an effective amount of an H₁ receptor antagonist selected from:
20 astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast,
25 pyrilamine, promethazine, terfenadine, tripelennamine, temelastine, trimeprazine or triprolidine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 287 in combination with an effective amount of an H₁ receptor antagonist selected from:
30 astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine,

descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast, pyrilamine, promethazine, terfenadine, tripeleennamine, temelastine, trimeprazine or
5 triprolidine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 320 in combination with an effective amount of an H₁ receptor antagonist selected from:
10 astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast,
15 pyrilamine, promethazine, terfenadine, tripeleennamine, temelastine, trimeprazine or triprolidine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 32 in combination with an effective amount of an H₁ receptor antagonist selected from:
20 loratadine, descarboethoxyloratadine, fexofenadine or cetirizine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 54 in combination with an effective amount of an H₁ receptor antagonist selected from:
25 loratadine, descarboethoxyloratadine, fexofenadine or cetirizine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 55 in combination with an effective amount of an H₁ receptor antagonist selected from:
30 loratadine, descarboethoxyloratadine, fexofenadine or cetirizine.

Another embodiment of this invention is directed to a method of treating: allergy, allergy-induced airway responses, and congestion comprising administering

- 20 -

to a patient in need of such treatment an effective amount of Compound 253A in combination with an effective amount of an H₁ receptor antagonist selected from: loratadine, descarboethoxyloratadine, fexofenadine or cetirizine.

Another embodiment of this invention is directed to a method of treating:
5 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 287 in combination with an effective amount of an H₁ receptor antagonist selected from: loratadine, descarboethoxyloratadine, fexofenadine or cetirizine.

Another embodiment of this invention is directed to a method of treating:
10 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 320 in combination with an effective amount of an H₁ receptor antagonist selected from: loratadine, descarboethoxyloratadine, fexofenadine or cetirizine.

Another embodiment of this invention is directed to a method of treating:
15 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 32 in combination with an effective amount of an H₁ receptor antagonist selected from: loratadine or descarboethoxyloratadine.

Another embodiment of this invention is directed to a method of treating:
20 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 54 in combination with an effective amount of an H₁ receptor antagonist selected from: loratadine or descarboethoxyloratadine.

Another embodiment of this invention is directed to a method of treating:
25 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 55 in combination with an effective amount of an H₁ receptor antagonist selected from: loratadine or descarboethoxyloratadine.

Another embodiment of this invention is directed to a method of treating:
30 allergy, allergy-induced airway responses, and congestion comprising administering to a patient in need of such treatment an effective amount of Compound 253A in combination with an effective amount of an H₁ receptor antagonist selected from: loratadine or descarboethoxyloratadine.

- 21 -


Another embodiment of this invention is directed to a method of treating:
allergy, allergy-induced airway responses, and congestion comprising administering
to a patient in need of such treatment an effective amount of Compound 287 in
combination with an effective amount of an H₁ receptor antagonist selected from:
5 loratadine or descarboethoxyloratadine.

Another embodiment of this invention is directed to a method of treating:
allergy, allergy-induced airway responses, and congestion comprising administering
to a patient in need of such treatment an effective amount of Compound 320 in
combination with an effective amount of an H₁ receptor antagonist selected from:
10 loratadine or descarboethoxyloratadine.


R¹ is preferably selected from:

- (A) aryl (most preferably phenyl);
(B) substituted aryl (e.g., substituted phenyl), wherein the
15 substituents on said substituted aryl are most preferably selected from: (1) halo (e.g.,
monohalo or dihalo), more preferably chloro or fluoro, even more preferably
monochloro, dichloro, monofluoro or difluoro; or (2) alkyl, more preferably unbranched
(i.e., straight chain, e.g., methyl) alkyl, even more preferably substituted alkyl, still
more preferably alkyl substituted with halo (e.g., 1, 2 or 3 halo atoms, such as Cl or
20 F), even still more preferably alkyl substituted with fluoro atoms, yet still more
preferably trifluoromethyl;

- (C) heteroaryl, most preferably a five or six membered heteroaryl
ring, more preferably a six membered heteroaryl ring, and still more preferably pyridyl,
examples of heteroaryl rings include pyridyl, thienyl, pyrimidinyl, thiazolyl or pyridyl N-
25 Oxide, most preferred heteroaryl rings are exemplified by




wherein




- 22 -

is preferred more;


- (D) substituted heteroaryl, most preferably halo or alkyl substituted heteroaryl (e.g., halopyridyl (e.g., fluoropyridyl) and alkylthiazolyl), more preferably substituted heteroaryl wherein the substituents are independently selected from the same or different alkyl groups (even more preferably one straight chain alkyl group, e.g., methyl), still more preferably alkyl substituted thiazolyl, and even more preferably



yet even more preferably



- (E) when R^1 is taken together with X, then the moiety is



wherein c is most preferably 0 or 1, and when c is 1 then R^6 is most preferably halo, and when c is 1 then R^6 is more preferably fluoro.

- X is preferably $=C(NOR^3)$ wherein R^3 is preferably selected from H, alkyl or halo substituted alkyl (e.g., fluoro substituted alkyl, such as $-\text{CH}_2\text{CF}_3$), most preferably alkyl, more preferably methyl or ethyl, and still more preferably methyl.

Preferably M^2 is nitrogen.

n is preferably 2.

a is preferably 0 or 1, and most preferably 0.

- b is preferably 0 or 1, and most preferably 0.

c is preferably 0 or 1, and most preferably 0, and when c is 1 then R^6 is preferably halo, and when c is 1 R^6 is most preferably fluoro.

e is preferably 1-5.


Y is preferably $=C(O)$ (i.e., $=C=O$).

- M^3 and M^4 are preferably selected such that: (1) one is carbon and the other is nitrogen, or (2) both are nitrogen, with M^3 most preferably being carbon.


- 23 -

p is preferably 2.


Z is preferably C₁ to C₃ alkyl, and most preferably



R² is preferably a six membered heteroaryl ring, most preferably pyridyl, substituted pyridyl, pyrimidinyl or substituted pyrimidinyl, more preferably pyridyl, pyridyl substituted with —NR⁴R⁵, pyrimidinyl or pyrimidinyl substituted with —NR⁴R⁵, still more preferably pyridyl, pyridyl substituted with —NH₂ (i.e., R⁴ and R⁵ are H), pyrimidinyl or pyrimidinyl substituted with —NH₂ (i.e., R⁴ and R⁵ are H), and even more preferably



and still even more preferably



R³ is preferably H or alkyl, most preferably H or methyl.

R⁴ is preferably H or lower alkyl, most preferably H or methyl, and more preferably H.

R⁵ is preferably H, C₁ to C₆alkyl or —C(O)R⁴, most preferably H or methyl, and more preferably H.

R¹² is preferably alkyl, hydroxyl or fluoro, and most preferably H.

R¹³ is preferably alkyl, hydroxyl or fluoro, and most preferably H.

Representative compounds of this invention include, but are not limited to: Compounds 23, 30, 31, 32, 33, 41, 44, 45, 49, 50, 52, 53, 54, 55, 56, 57A, 59, 65, 75, 76, 80, 82, 83, 88, 92, 99, 104, 105, 110, 111, 117, 121, 123, 127, 128, 200-241, 244-273, 275, and 278-282, 287, 296, 301-439 and 446.

Thus, representative compounds of this invention include, but are not limited to: Compounds 23, 30, 31, 32, 33, 44, 45, 49, 50, 53, 54, 55, 59, 75, 76, 83, 88, 92,

- 24 -

99, 104, 110, 117, 128, 200, 201, 203-215, 217-241, 244-246, 246A, 247-253, 253A, 254-273, 275, 278, and 280-282, 317, 334 and 403.

Preferred compounds of this invention are selected from: Compound 23, 30, 31, 32, 33, 50, 53, 54, 55, 56, 57A, 59, 92, 212, 215, 218, 219, 220, 224, 225, 226, 227, 229, 233, 235, 237, 238, 246, 246A, 247, 248, 251, 253, 253A, 268-273, 275, 278-281, 287, 296, 301, 304-307, 309, 312, 314-318, 320-356, or 358-376.

Most preferred compounds of this invention are selected from: Compound 30, 31, 32, 33, 54, 55, 56, 57A, 225, 237, 246A, 253A, 273, 280, 287, 296, 301, 304-307, 309, 312, 314-318, 320-348, 350-356, 359-372, and 374-376.

Thus, one embodiment of this invention is directed to Compound 32.

Another embodiment of this invention is directed to Compound 54.

Another embodiment of this invention is directed to Compound 55.


Another embodiment of this invention is directed to Compound 253A.

Another embodiment of this invention is directed to Compound 287.


Another embodiment of this invention is directed to Compound 320.

Structures for the above compounds are found in the Examples below, and in Tables 1 to 3 below.

The more preferred compound of this invention is the compound of the formula:




This invention also provides a compound of the formula:



This invention also provides a compound of the formula:


- 25 -




Compounds 32A and 32B can also be used in the pharmaceutical compositions, and the methods of this invention.

The following processes may be employed to produce compounds of the invention.

One synthetic route involves a linear sequence of reactions to obtain the desired compounds, i.e.,




This linear sequence of reactions to synthesize compounds of this invention is illustrated below. In the illustrated procedure R¹ is aryl, heteroaryl, or alkyl; X = a ketone, oxime or substituted oxime; M¹ = M³ = carbon; M² = M⁴ = nitrogen; Y is C=O; Z = CHR; R² is heteroaryl; and n and m = 2 (n and m being 1 can also be prepared by this procedure).

Step 1 Synthesis of Ketone 8

In the above equations PG represents a protecting group, and M represents Li or MgX^1 (wherein X^1 represents Cl, Br or I).


In equation 1 and 2, a Grignard reagent 2 is reacted with an electrophile such as the aldehyde 1 or the nitrile 4 in a suitable aprotic solvent such as THF or ether. PG represents a protecting group. Suitable protecting groups include, for example, methyl and benzyl. In the case of nitrile 4, acidic workup yields the ketone 8 directly. Alcohol 3 can be oxidized by a number of different reagents to give 8. Alternatively, the amide 7 can be reacted with an organometallic reagent to directly give the ketone 8. Suitable protecting groups for this step include carbamates or amides or the like. Thus, examples of protecting groups in equation 3 include t-BOC, CBZ and FMOC.

Step 2: Deprotection of 8

- 27 -


When the protecting group, PG, is a methyl group, said methyl group can be removed using a reagent such as a chloroformate; when PG is a carbamate, such as, a t-Boc group, it can be removed by dilute acid, such as, for example HCl.

5 Step 3: Synthesis of 11




Amine **9** can be coupled to acid **10** using a number of methods well known in the art such as DCC or PyBOP. Alternatively, the acid **10** can be activated by conversion to the acid chloride or mixed anhydride and then reacted with the amine **9** to give **11**. Suitable protecting groups for **10** include, for example, t-Boc.

Step 4: Synthesis of Amine 12



Compound **11** in which the protecting group is a t-Boc can be deprotected under acidic conditions such as HCl in dioxane or TFA in CH₂Cl₂ to give the amine **12**.

Step 5: Synthesis of Compound 14




R^{30} in **13** represents an alkyl group. E is a leaving group, halogen, or E is a carbonyl group.

Compound **14** can be prepared by reacting amine **12** with **13**. When E represents a carbonyl group (C=O), **12** and **13** are combined in a solvent such as

- 28 -




CH₂Cl₂ in the presence of molecular sieves. After the reaction is complete (e.g., 1 to 10 h), a reducing agent such as NaBH(OAc)₃ is added. Alternatively, when E is a halogen atom such as Cl or Br, 12 and 13 are combined in a solvent, such as DMF, in the presence of a tertiary amine base to give the product 14. Suitable protecting groups include, for example t-Boc, phthaloyl.

Step 6: Synthesis of Compound 16




Compound 14 can be converted to the oxime 15 by combining 14 with H₂NOR³·HCl in pyridine at a temperature of 40 – 60°C. Alternatively, 14 can be combined with H₂NOR³·HCl in an alcoholic solvent in the presence of a base, such as, NaOAc, to give 15.

An alternate approach to the synthesis of compounds of Formula I involves the synthesis of the two halves of the molecule followed by coupling of the two pieces, i.e.,



In this case, the synthesis of the AB fragment is the same as that described above. The synthesis of the CD fragment is given below.

Step 1: Synthesis of Compound 17




R³⁰ is as defined above (i.e., alkyl). R³⁵ is methyl or ethyl.

- 29 -

Compound 17 is synthesized in the same manner as that described for the synthesis of compound 14.

Step 2: Synthesis of Compound 18



M represents Li, Na, or K.


Compound 17 is saponified in a mixed solvent, such as, for example: (1) EtOH or MeOH and water, or (2) THF, water, and MeOH, using an alkali metal base such as LiOH or NaOH at a temperature of from 50 to 100°C to give the salt 18.

Compound 18 can be combined with compound 9, as described above, to give 14. The remaining steps are the same.

Compounds useful in this invention are exemplified by the following examples which should not be construed as limiting the scope of the disclosure. Alternative mechanistic pathways and analogous structures within the scope of the invention may be apparent to those skilled in the art.


Example 1

Step 1




To a solution of 10.81 g (100 mmol) of 2-amino-4-methylpyridine in 250 ml of tert-butanol was added 26.19 g (120 mmol) of BOC anhydride. The reaction mixture was stirred at room temperature overnight, concentrated, loaded on silica gel and flash chromatographed (from 30% hexanes/ CH₂Cl₂ to 0 - 2% acetone/ CH₂Cl₂) to produce 15.25 g (73.32 mmol; 73%) of 1A as a white solid.

- 30 -

Step 2

To a -78°C solution of 1A (35.96 g, 173 mmol) in THF (1.4 L) was added 1.4 M BuLi solution (272 ml, 381 mmol) in hexanes in portions over 30 min. Reaction mixture was then allowed to warm up and was stirred for 2 h at room temperature, which resulted in the formation of an orange precipitate. The mixture was cooled back to -78°C , and predried oxygen (passed through a Drierite column) was bubbled through the suspension for 6 h while the temperature was maintained at -78°C . Reaction mixture color changed to yellow during this time. It was then quenched at -78°C with 51.4 ml (700 mmol) of Me_2S followed by 22 ml (384 mmol) of AcOH. Reaction mixture was allowed to warm up and was stirred for 48 h at room temperature. Dilution with water and extraction with EtOAc were followed by concentration and flash chromatography (0 – 15% acetone/ CH_2Cl_2) to provide 20.15 g (90 mmol; 52%) of alcohol 2A as a pale yellow solid.


Step 3

To a solution of 19.15 g (85.5 mmol) of alcohol 2A in 640 ml of CH_2Cl_2 was added saturated aqueous solution of 8.62 g (103 mmol) of NaHCO_3 and 444 mg (4.3 mmol) of NaBr. Reaction mixture was cooled to 0°C , and 140 mg (0.90 mmol) of TEMPO was introduced. Upon vigorous stirring 122 ml of 0.7 M (85.4 mmol) commercial bleach solution (5.25% in NaOCl) was added in portions over 40 min. After additional 20 min at 0°C reaction mixture was quenched with saturated aqueous $\text{Na}_2\text{S}_2\text{O}_3$ and allowed to warm to room temperature. Dilution with water and

- 31 -

extraction with CH_2Cl_2 were followed by concentration and flash chromatography (from 30% hexanes/ CH_2Cl_2 to 0 - 2% acetone/ CH_2Cl_2) to afford 15.97 g (71.9 mmol; 84%) of aldehyde **3A** as an off-white solid.


5 **Step 4**



To a solution of 11.87 g (53.5 mmol) of aldehyde **3A** in 370 ml of CH_2Cl_2 was added 9.07 ml (58.8 mmol) of ethyl isonipecotate followed by four drops of AcOH. Reaction mixture was then stirred for 40 min at room temperature after which 22.68 g (107 mmol) of $\text{NaBH}(\text{OAc})_3$ was introduced. Reaction mixture was stirred overnight at room temperature, neutralized with saturated aqueous NaHCO_3 , diluted with water and extracted with CH_2Cl_2 . Concentration and flash chromatography (0 – 4% sat. NH_3 in MeOH/ CH_2Cl_2) provided 19.09 mg (52.6 mmol; 98%) of **4A** as an off-white solid.


15

Step 5




To a solution of 1.57 g (4.33 mmol) of ester **4A** in 10 ml of a 3 : 1 : 1 mixture of THF – water – methanol was added 0.125 g (5.21 mmol) of LiOH. Reaction mixture was stirred overnight at room temperature, concentrated and exposed to high vacuum to obtain 1.59 g of crude acid **5A** as a yellowish solid which was used without purification.

- 32 -


Example 2

A solution of compound **6A** (42mmol), NBS (126mmol) and Bz₂O₂ (4.2mmol) in CCl₄ (400ml) was refluxed at 80°C for 5 h, cooled and stirred at room temperature overnight. The reaction was filtered and concentrated, and the residue was purified by flash column (30% EtOAc/Hexane) to obtain the desired compound **7A** (3.1g, 23%).

Example 3**Step 1**


To a solution of **8A** (10 g, 79.4 mmol) and DMAP (0.029 g, 0.24 mmol) in methylene chloride (150 mL) at 0°C was added phthaloyl dichloride (16.1 g, 79.4 mmol) dropwise. The reaction mixture was stirred at room temperature overnight. After stirring overnight, the reaction was washed with saturated aqueous NaHCO₃, water, dried and concentrated to give compound **9A** as a yellow solid (20 g, 99.8%) which was used without further purification.

- 33 -

Step 2


In a manner similar to that described in Example 2, compound **9A** (20 g, 79.3 mmol) was converted to compound **10A**.

5

Step 3

Compound **10A** (0.5 g, 1.5 mmol) and hydrazine (0.5 M in ethanol, 5 mL, 2.5 mmol) were combined and stirred at room temperature overnight. The reaction was diluted with water and extracted with methylene chloride. The organic layer was dried, concentrated and the residue purified on a flash column (3% methanol in ethyl acetate) to give compound **11A** (0.2 g, 66%).

10

Example 4**Step 1**

15

- 34 -

Compounds **12A** (2 g, 18.3 mmol) and **13A** (3.5 g, 22 mmol) were dissolved in methylene chloride and stirred at room temperature for 1 h. NaB(OAc)₃H (5.4 g, 25.6 mmol) was added and the mixture stirred at room temperature for 5h. The reaction was washed with saturated aqueous NaHCO₃, dried and concentrated, and the residue purified by flash column (2% methanol in ethyl acetate). Compound **14A** was obtained (4.5g, 99%).


Step 2



In a manner similar to that described in Example 1, Step 5, compound **14A** (0.35 g, 1.4 mmol) was converted to compound **15A** (0.31g, 100%).


Example 5

Step 1




To the solution of 2,4-difluorobenzaldehyde (**16A**, 28.1mmol) in THF (10ml) was added the Grignard reagent **17A** (1.33M in THF, 30ml), and the mixture was stirred at room temperature overnight. The reaction was quenched with saturated NH₄Cl (150ml), extracted three times with EtOAc (100ml), dried, filtered and concentrated. Flash chromatography (20% MeOH/EtOAc) yielded the desired compound **18A** (1.8g, 27%).


- 35 -

Step 2

Compound **18A** (1.6 g, 6.7 mmol), $\text{H}_2\text{NHOH}\cdot\text{HCl}$ (0.95 g, 6.7 mmol) and pyridine (10 mL) were combined and heated to 60°C overnight. The pyridine was removed under vacuum and the residue treated with methylene chloride and saturated aqueous NaHCO_3 . The organic layer was separated, dried, and concentrated, and the residue purified by flash chromatography to give compound **19A** (1.4g, 82%).


Step 3

To the suspension of NaH (0.41g, 10.2 mmol) in THF (10ml) was slowly added a solution of **19A** (1.3 g, 5.11 mmol) in DMF (5 mL) dropwise and the reaction stirred at 70~75°C overnight. The mixture was extracted twice with EtOAc and three times with H_2O (30 mL), dried over MgSO_4 and concentrated to give crude **20A** which was used without further purification (1.04 g, 87%).

Step 4


To the solution of compound **20A** (4.3 mmol) in dichloroethane (20 ml) at 0°C was added 2-chloroethyl chloroformate (6.2 mmol) and triethylamine (7.2 mmol) and the reaction was stirred at room temperature overnight. The solvent was evaporated, Et₂O was added to the residue, and the unreacted starting material was removed by filtration. The filtrate was concentrated and the residue redissolved in MeOH and refluxed for 30 min. Removal of the methanol gave the product **21** (0.3g) which was used without further purification.

Step 5



To a mixture of compound **21** (1.64 mmol), compound **5A** (1.64 mmol) and PyBOP (1.64 mmol) was added DIPEA (4.92 mmol) and CH₂Cl₂ (10 ml), and the reaction was stirred over the weekend at room temperature. Saturated NaHCO₃ (100 ml) was added and the reaction was extracted and twice with CH₂Cl₂ (100 mL), dried over solid MgSO₄, concentrated and flash chromatographed (70% EtOAc/Hexane) to give compound **22** (1.04 mmol, 64%).

Step 6




Compound **22** (0.2 g, 0.37 mmol) was dissolved in CF₃CO₂H (3 mL) and methylene chloride (3 mL) and stirred at room temperature overnight. The solvent was removed by evaporation, saturated aqueous NaHCO₃ was added and mixture

extracted with methylene chloride. The organic layer was dried (MgSO_4), filtered and concentrated, and the residue purified by flash chromatography to give compound **23** (0.11 g, 68%).


5 Example 6

Step 1




A solution of **24** (50 g, 387 mmol) and triethylamine (110 mL) in dioxane (400 mL) and water (400 mL) at 4°C was treated with Boc₂O (93 g, 426 mmol). The cooling bath was removed and the solution allowed to warm to room temperature. After 21h, the volume was reduced by two-thirds under vacuum. The residue was poured into ethyl acetate (250 mL) and water (250 mL). Saturated aqueous NaHCO₃ (250 mL) was added and the organic phase was separated and discarded. The aqueous phase was acidified with 10% HCl and extracted with ethyl acetate. The combined organic phases were washed with water, brine, and dried (Na₂SO₄), and concentrated to give **25** as a white powder (82 g, 94%).

Step 2




To a solution of compound **25** (40 g, 175 mmol) in DMF (250 mL) at 4°C was added N,O – dimethylhydroxylamine, hydrochloride (34 g), EDCI (44 g, 0.228 mol), HOBT (2.4 g), and DIPEA (120 mL). The reaction was warmed to room temperature and stirred overnight. The reaction was then concentrated to half volume in vacuo and poured onto 1:1 ethyl acetate:water. The organic layer was separated and the aqueous layer extracted with additional ethyl acetate. The combined organic layers were washed with saturated aqueous NH₄Cl, saturated aqueous NaHCO₃, water, and brine, and dried. Concentration gave **26** as a light yellow oil (46.7 g, 99%)

- 38 -


Step 3

To a solution of 2-bromopyridine (17.6 mL, 0.184 mol) in THF (600 mL) at -78°C was added n-BuLi (115 mL of a 1.6M solution in hexanes, 0.184 mol) dropwise over 15 min. After stirring for an additional 30 min at this temperature, a solution of 26 (25 g, 91.9 mmol) in THF (500 mL) was added dropwise over 15 min. The reaction was removed from the cold bath and placed in an oil bath and heated to 60°C for 1.5h. The reaction was then cooled to 4°C , diluted with ether (500 mL), and treated with saturated aqueous Na_2SO_4 (≈ 5 mL). The mixture was transferred to an Erlenmeyer flask and diluted with additional ether (700 mL). Additional saturated aqueous Na_2SO_4 was added followed by solid Na_2SO_4 . The mixture was filtered through a plug of solid Na_2SO_4 and concentrated in vacuo. Flash column chromatography (0-20% ethyl acetate in hexanes) yielded compound 27 as a yellow oil (16.85 g, 63%).


Step 4

A solution of 27 (3.3 g, 11.4 mmol) in methanol (50 mL) was treated with 4M HCl in dioxane (50 mL) and stirred at room temperature for 1.5 h. Removal of the solvent in vacuo gave 28 as a tan powder (3g, 100%).

- 39 -

Step 5


To a suspension of compound 5A (17.4 g, 50 mmol), compound 28 (11 g, 42 mmol), and diisopropylethylamine (34.6 mL, 199 mmol) in DMF (125 mL) was added
 5 HOBT (7.83 g, 58 mmol), EDC (18.54 g, 96.7 mmol), and 4Å molecular sieves. The mixture was stirred for 40 h at room temperature, diluted with methylene chloride (600 mL) and 0.5 N NaOH (400 mL) and filtered. The precipitate was washed thoroughly with additional 0.5N NaOH and methylene chloride. The combined organic phases
 10 were concentrated and chromatographed twice on silica gel (1:1 hexane:methylene chloride to 6% saturated NH₃ in methanol in methylene chloride) to produce 29 as a tan solid (22.3 g) which was used as is in the next step.

Step 6

15 A solution of 29 (22.3 g, 44 mmol) in methylene chloride (120 mL) and trifluoroacetic acid (60 mL) was stirred for 7 h at room temperature. The reaction was


- 40 -

concentrated, exposed to high vacuum for 3h, dissolved in toluene and concentrated and then exposed again to high vacuum. The so-obtained crude brown oil was used in the next step without further purification.

5 **Step 7**


Compound 30 (≈ 17.9 g, 44 mmol) was dissolved in pyridine (420 mL), treated with $\text{H}_2\text{NOCH}_3 \cdot \text{HCl}$ (21.78 g, 264 mmol) and heated to 90°C for 14h. The reaction was then concentrated and the residue taken up in a mixture of methylene chloride, (500 mL) and 2N NaOH (500 mL). The organic phase was separated and the aqueous phase extracted with additional methylene chloride (300 mL). The organic phases were dried and concentrated, and the residue chromatographed on SiO_2 (0-13% NH_3/MeOH in CH_2Cl_2) to produce a yellow solid (9.26 g). The mixed fractions from the column were rechromatographed to give an additional 3.23g of the desired material. Total yield 12.49 g (65% yield over the last two steps).

- 41 -

Step 8


Compound **31** (1 g) in ethanol (15 mL) was separated into the pure isomers using a Chiralcel AD column (20 mm x 500 mm) (eluent: 75:25 hexane: isopropanol plus 0.5% N,N-diethylamine; flow rate: 50 mL/min; UV detection at 254 nm) to give compound **32** (0.6 g) and compound **33** (0.4 g). $[M+H]^+$ 437 for **32** and **33**.

Alternatively, compound **32** can preferably be prepared from compound **5A** in a manner similar to that described for compound **287** in Step 3 of Example 28.


Example 7**Step 1**

To a solution of **34** (2.4 g, 13.5 mmol) in THF (15 mL) was added compound **35** (26 mL of a 1.3M solution) and the reaction stirred overnight at room temperature. 2N HCl was then added till the pH < 2 and the THF was removed under reduced pressure. The pH was neutralized by the addition of 1N NaOH and the aqueous phase extracted with 5% MeOH in EtOAc. The organic phase was dried, concentrated, and the residue chromatographed (20% MeOH in EtOAc) to give **36** (1.03 g, 28%).


- 42 -

Step 2

To a solution of **36** (1.03 g, 3.78 mmol) in 1,2-dichloroethane (30 mL) was added 1-chloroethylchloro formate (0.76 mL, 7.6 mmol) and the reaction stirred at room temperature overnight. The solvent was removed in vacuo and the residue washed with ether. Solid residue was removed by filtration and the ether removed by evaporation to give an oil which was dissolved in MeOH (15 mL) and heated to reflux for 2h. Removal of the solvent gave **37** which was used in the next step without further purification (1.4 g).


Step 3

Compound **37** (0.98 g, 3.78 mmol), N-Boc isonipocotic acid (0.87 g, 3.78 mmol), DEC (1.11 g, 5.7 mmol), HOBT (0.68g, 4.91 mmol) and DIPEA (3 mL) were combined in CH_2Cl_2 (40 mL) and stirred overnight at room temperature. The reaction was then diluted with CH_2Cl_2 and washed with saturated aqueous NaHCO_3 . The organic layer was dried, concentrated and the residue chromatographed (10% hexane in EtOAc) to give **38** (1.61 g, 91%).

Step 4


- 43 -

Compound **38** (1.61 g, 3.43 mmol) in CH_2Cl_2 (15 mL) was treated with 1N HCl in dioxane (5.2 mL) and stirred overnight at room temperature. The solvent was removed in vacuo to give **39** (1.65 g) which was used without further purification.


5 **Step 5**

Compound **39** (1.65 g, 4.01 mmol), **7** (1.29 g, 4.07 mmol) and Et_3N (1.7 mL) were combined in DMF (40 mL) and stirred at room temperature overnight. The reaction was dissolved in EtOAc and washed 4 times with water. The organic layer was dried and concentrated, and the residue purified by chromatography (5% MeOH in EtOAc) to give **40** (0.6 g, 47%).

- 44 -


Step 6

A solution of 40 (0.31 g, 0.51 mmol) in pyridine (5 mL) was treated with $\text{H}_2\text{NOMe}\cdot\text{HCl}$ (0.092 g, 1.08 mmol) and heated to 60°C overnight. The reaction was
5 diluted with 10% MeOH in CH_2Cl_2 , washed with saturated aqueous NaHCO_3 , dried, and concentrated, and the residue purified by chromatography (10–15% MeOH in EtOAc) to give 41 (0.09 g).


Example 810 **Step 1**

In a manner similar to that described in Example 7, Steps 3–4, compound 42 was converted to compound 43.


- 45 -

Step 2

To a solution of **43** (2.3 g, 6.3 mmol) in CH_2Cl_2 (60 mL) was added 4Å molecular sieves and 4-formylpyridine (0.68 mL, 6.9 mmol) and the mixture stirred for 3 h at room temperature. $\text{Na}(\text{OAc})_3\text{BH}$ (2.7 g, 12.7 mmol) was then added and the reaction stirred for 1h. The reaction was quenched by the addition of NH_4Cl followed by the addition of saturated aqueous NaHCO_3 . The reaction mixture was then extracted with EtOAc, and the combined organic layers were dried and concentrated to give a residue which was chromatographed (20% MeOH in EtOAc). Compound **44** was obtained (2.3 g, 87%).

Step 3


In a manner similar to that described in Example 7, Step 6, compound **44** was converted to compound **45**.

Example 9.**Step 1**

- 46 -


In a manner similar to that described in Example 8, Step 2, compound 46 (1.13 g, 6 mmol) was converted to compound 47 (1.7 g, 100%).

Step 2



In a manner similar to that described in Example 7, Step 4, compound 47 (1.7 g, 6.13 mmol) was converted to compound 48 (1.9 g, 100%).


Step 3



A mixture of compound 48 (0.57 g, 2 mmol) and compound 42 (0.52 g, 2 mmol) in CH₂Cl (20 mL) was added Et₃N (1.95 mL) and the reaction cooled to -40°C. Triphosgene (0.2 g) was added and the reaction stirred at -40°C for 2 h and room temperature for 48 h. The reaction was then washed with 1N NaOH, brine, and the organic layer dried. Concentration gave a residue that was purified by column chromatography (10% MeOH in EtOAc) to give 49 (0.14 g, 55%).


15

Step 4




- 47 -


In a manner similar to that described in Example 7, Step 6, compound **49** (0.09 g, 0.21 mmol) was converted to compound **50**.

Example 10**Step 1**

In a manner similar to that described in Example 7, Steps 3-4, compound **28** (2.6 g, 9.9 mmol) was converted to compound **51** (1.1 g).



Step 2

In a manner similar to that described in Example 7, Step 5, compound **51** (1.1 g, 2.94 mmol) was reacted with compound **11** (0.59 g, 2.94 mmol) to give compound **52** (0.53 g).



Step 3

In a manner similar to that described in Example 6, Step 7, compound **52** (0.53 g, 1.26 mmol) was converted to compound **53** (0.48 g).

- 48 -

Step 4

5



In a manner similar to that described in Example 6, Step 8, the 4 diastereomers
10 of compound 53 could be obtained using a Chiralcel AD column (75:25 hexane:EtOAc
plus 0.5% Et₂NH). The two faster eluting compounds (54 and 55) were the E-oxime
isomers and the slower eluting compounds (56 and 57A) were the Z-oxime isomers.

Isomer A	54	0.12 g
Isomer B	55	0.11 g
Isomer C	56	0.08 g
Isomer D	57A	0.06 g


Example 11

Step 1




A solution of n-BuLi (4.2 mL of a 1.6 M solution in hexane) in THF (25 mL) was treated at -25°C with (i-Pr) $_2$ NH (0.69 g, 6.8 mmol). The reaction was stirred for 1 h at 0°C and then cooled to -70°C . Compound **4A** (0.82 g, 2.26 mmol) in THF (5 mL) was added dropwise and the reaction stirred at -70°C for 2 h and -50°C for 2 h. The reaction was recooled to -70°C and (1S)-(+)-(10-camphorsulfonyl)oxaziridine (1.04 g, 4.52 mmol) in THF (5 mL) was added. The reaction was stirred at -70°C for 2 h and slowly warmed to room temperature overnight. The reaction was quenched by the addition of saturated aqueous NH_4Cl and extracted with EtOAc. The organic layer was dried and concentrated, and the residue purified by column chromatography (1:1 hexane:EtOAc) to give **57** (0.44 g, 51%).

Step 2




In a manner similar to that described in Example 1, Step 5, compound 57
20 (0.42 g, 1.1 mmol) was converted to compound 58 (0.4 g).

- 50 -

Step 3


In a manner similar to that described in Example 6, Steps 5 - 8, compound 58 (0.25 g, 0.7 mmol) was converted to compound 59 (0.1 g).

5

Example 12**Step 1**

A solution of compound 60 (10 g, 50.7 mmol) in ether (150 mL) at -78°C was treated sequentially with TMEDA (11.8 g, 101.4 mmol) and *s*-BuLi (58.5 mL of a 1.3M solution in hexanes, 76 mmol) and the reaction stirred at this temperature for 6 h. Neat $\text{CH}_3\text{SO}_4\text{CH}_3$ (12.8 g, 101.4 mmol) was then added and the reaction allowed to slowly warm to room temperature overnight. Saturated aqueous NaCl was added and the organic layer was separated. The aqueous layer was extracted three times with ether and the combined organic layers were dried, concentrated, and the residue chromatographed (5% EtOAc in hexane) to give 61 (8.0 g, 75%).

15


Step 2

20


A solution of 61 (8 g, 37.9 mmol) in THF (40 mL) at 0°C was treated dropwise with a solution of $\text{BH}_3 \cdot \text{THF}$ (45.4 mL of a 1.0M solution in THF, 45.4 mmol) and the

- 51 -

reaction allowed to slowly warm to room temperature overnight. The reaction was recooled to 0°C, EtOH (13 mL), pH = 7 buffer (25 mL) and H₂O₂ (25 mL) was added, and the reaction allowed to stir at room temperature overnight. The solvent was then removed in vacuo and the residue poured into water and CH₂Cl₂. 10% aqueous NaOH (10 mL) was added and the organic layer separated. The aqueous layer was extracted with additional CH₂Cl₂ and the combined organic layers were dried and concentrated. The residue was chromatographed (40% EtOAc in hexane) to give 62 (3 g).

10 **Step 3**


A solution of 62 (2.8 g, 12.2 mmol) in EtOAc (30 mL) and NaBr (1.26 g, 0.12 mmol) in saturated aqueous NaHCO₃ (30 mL) was cooled to 0°C and treated with TEMPO (0.02 g, 0.12 mmol). After 15 min., NaOCl (17.44 mL) was added and the mixture stirred for 3 h. Saturated aqueous Na₂S₂O₃ was added and the pH adjusted to 5 – 6 by the addition of 1N HCl. The mixture was extracted with EtOAc and the organic layers were dried and concentrated. The residue was chromatographed (10 – 20% EtOAc in hexane) to give compound 63 (2.1 g, 76%).

20 **Step 4**


To a cooled (0°C) suspension of PCC (0.95 g, 4.4 mmol) in CH₂Cl₂ (5 mL) was added dropwise a solution of 63 (0.5 g, 2.2 mmol). And the mixture stirred overnight at room temperature. Additional PCC (1 eq.) was added and the mixture was heated

- 52 -

to reflux for 2 h. The reaction was cooled, filtered through celite, and concentrated to give crude **64** (1.5 g) which was used without further purification.


Step 5

In a manner similar to that described in Example 5, Step 5, Example 7, Step 4, Example 1, Step 4, and Example 6, Steps 6 and 7, **64** (0.73 g, 3 mmol) was converted to **65** (0.1 g).


10 **Example 13****Step 1**

To a 0°C solution of Vilsmeier salt, prepared by the dropwise addition of phosphorus oxychloride (150.0 mL; 1.61 mol) to DMF (310.4 mL; 4.01 mol) over 15 min. and subsequent cooling in an ice bath, was added malonic acid (40.1 g; 0.39 mol) in portions over 45 min. The reaction mixture was then heated to 100°C, and the stirring was continued for 48 h. The reaction mixture was then allowed to cool to room temperature and was quenched by slowly pouring it into a suspension of NaHCO₃ (808 g; 9.62 mol) in water. The solution was decanted off the excess of NaHCO₃ and concentrated to dryness under vacuum. After exposure to high vacuum for 2 days, the solid residue was washed repeatedly with CH₂Cl₂ until TLC indicated complete removal of product. Combined organic extracts were concentrated under vacuum to produce 41.0 g of dark brown oil, which was used directly in the next step.


- 53 -

Step 2

To a solution of 32.5 g (256 mmol) of crude malondialdehyde 66 in 650 ml of absolute ethanol was added 24.5 g (256 mmol) of guanidine hydrochloride and 17.4 g (256 mmol) of sodium ethoxide. The reaction mixture was refluxed for 4 h, cooled down to room temperature, concentrated and dry loaded on silica gel under vacuum. Flash chromatography (0-10% MeOH/ 20% acetone/ CH₂Cl₂) afforded 11.0 g (89.4 mmol; 23% from malonic acid (2 steps)) of pyrimidine 67 as a light yellow solid.


Step 3

To a mixture of 166 mg (1.35 mmol) of aminopyrimidine 67, 17 mg (0.14 mmol) of DMAP and 418 μ L (3.00 mmol) of Et₃N in 10 mL of THF was added 589 mg (2.7 mmol) of (BOC)₂O. The mixture was stirred at room temperature for 5 h, concentrated-dry loaded on silica gel and flash chromatographed (1-3% acetone/ CH₂Cl₂) to produce 117 mg (0.36 mmol; 27%) of 68 as a clear oil.


Step 4

- 54 -

To a solution of 117 mg (0.36 mmol) of aldehyde 68 in 7 mL of CH₂Cl₂ was added 67 μ L (0.43 mmol) of ethyl isonipecotate and 5 μ L of acetic acid. 30 min. later 153 mg (0.72 mmol) of NaBH(OAc)₃ was introduced. The mixture was stirred overnight at room temperature, diluted with CH₂Cl₂, washed with aqueous NaHCO₃, dried and concentrated, and crude residue was flash chromatographed (0-4% sat. NH₃ in MeOH/ CH₂Cl₂) to produce 133 mg (0.29 mmol; 81%) of 69 as a white film.

Step 5

To a solution of ester 69 in 5 mL of a 3 : 1 : 1 mixture of THF – water – methanol was added 11 mg (0.44 mmol) of LiOH. Reaction mixture was stirred overnight at room temperature, concentrated to dryness and exposed to high vacuum to obtain 134 mg of crude acid 70 as a yellowish solid which was used without purification.

Example 14**Step 1**

To a -78°C solution of 2.36 g (11.4 mmol) of picoline 1A in 70 mL of THF was added 16.3 mL of 1.4 M BuLi solution (22.8 mmol) in hexanes in portions over 10 min.


- 55 -

Reaction mixture was then allowed to warm up and was then stirred for 2 h at room temperature, which resulted in the formation of an orange precipitate. The mixture was cooled back to -78°C , and ethylene oxide was bubbled through the solution for 1 min. followed by stirring for 5 min. This two-step sequence was repeated eight times.

5 The mixture was then allowed to warm to -50°C , stirred at that temperature for 40 min., quenched with 1.34 mL (23 mmol) of AcOH and allowed to warm to room temperature. Dilution with water was followed by extraction with EtOAc, concentration of the organic phase, and flash chromatography of the crude residue (10-15% acetone/ CH_2Cl_2) to produce 1.50 g (5.95 mmol; 53%) of 71 as a white solid.

10

Step 2




To a -60°C solution of 628 μL (7.2 mmol) of oxalyl chloride in 20 mL of CH_2Cl_2 was added dropwise 1.03 mL (14.5 mmol) of DMSO. After stirring the mixture for 15 min. at -55°C , a solution of 1.50 g (5.95 mmol) of alcohol 71 in 20 mL of CH_2Cl_2 was introduced over the period of 15 min. After the addition was complete, the mixture was stirred for 30 min. at -55°C , followed by the addition of 4.18 mL (30.0 mmol) of Et_3N and stirring for another 15 min. The reaction mixture was then warmed to room temperature and diluted with water. Extraction with CH_2Cl_2 was followed by


15

20 concentration of the organic phase and flash chromatography (1-15% acetone/ CH_2Cl_2) to produce 1.00 g (4.00 mmol; 67%) of 72 as an off-white solid.

- 56 -


Step 3

To a solution of 1.00 g (4.0 mmol) of aldehyde **72** in 25 mL of CH_2Cl_2 was added 617 μL (4.8 mmol) of ethyl isonipecotate followed by one drop of AcOH. Reaction mixture was then stirred for 40 min at room temperature after which 1.70 g (8.0 mmol) of $\text{NaBH}(\text{OAc})_3$ was introduced. Reaction mixture was stirred overnight at room temperature, neutralized with saturated aqueous NaHCO_3 , diluted with water and extracted with CH_2Cl_2 . Concentration and flash chromatography (0 – 4% saturated NH_3 in $\text{MeOH}/\text{CH}_2\text{Cl}_2$) provided 1.41 g (3.6 mmol; 90%) of **73** as a white solid.

Step 4


To a solution of 534 mg (1.47 mmol) of ester **73** in 4 mL of a 3 : 1 : 1 mixture of THF – water – methanol was added 60 mg (2.50 mmol) of LiOH. Reaction mixture was stirred overnight at room temperature, concentrated to dryness and exposed to high vacuum to obtain 540 mg of crude acid **74** as a white solid which was used without purification.

- 57 -

Example 15


In a manner similar to that described in Example 6, steps 5, 6, and 7, **70** was converted to **75**.

5

Example 16

In a manner similar to that described in Example 6, steps 5, 6, and 7, **74** was converted to **76**.

10


Example 17**Step 1**

To a solution of **77** (0.73 g, 3.82 mmol) in CH_2Cl_2 (10 mL) was added $(\text{COCl})_2$ (0.41 mL, 4.58 mmol) followed by DMF (0.1 mL) and the reaction was maintained at 40°C for 3 h. The reaction was then concentrated to give a brown solid which was dissolved in CH_2Cl_2 (10 mL). N , O -dimethylhydroxylamine hydrochloride (0.56 g, 5.73


15

- 58 -

mmol) and DIPEA (1.33 mL) were added and the reaction was stirred at room temperature overnight. The reaction was quenched by the addition of saturated aqueous NaHCO₃ and extracted with EtOAc. The combined organic layers were dried and concentrated, and the residue purified by chromatography to give **78** (3.2 g, 84%).


Step 2

In a manner similar to that described in Example 5, steps 1 and 4, **78** (0.57 g, 2.41 mmol) was converted to **79** (0.59 g).


Step 3

In a manner similar to that described in Example 6, steps 5, 6 and 7, **79** (0.38 g, 1.49 mmol) was converted to **80** (0.24 g).

- 59 -


Example 18**Step 1**

In a manner similar to that described in Example 6, step 7, **81** (0.36 g, 0.53 mmol; synthesized in the same manner as compound **30**) was converted to **82** (0.34 g, 63%).


Step 2

To a solution of **82** (0.115 g, 0.25 mmol) in DMF (4 mL) was added NaH (60% dispersion in mineral oil, 0.03g, 0.76 mmol). After 5 h at room temperature, $\text{CF}_3\text{CH}_2\text{OSO}_2\text{CF}_3$ (0.069 g, 0.3 mmol) was added and the reaction stirred at room temperature overnight. The reaction was diluted with EtOAc and extracted 3 times with water to remove the DMF. The organic layer was dried and concentrated to give a residue which was purified by chromatography (10% MeOH/ NH_3 in EtOAc) to give **83** (0.08 g, 30%).


- 60 -

Example 19**Step 1**

To a solution of 17 (0.21 mole, 100ml THF, -10°C) was added 84 (0.14 mole) over 5 min and the reaction mixture became very viscous. Additional THF (100ml) was added and the yellow suspension was warmed from -10°C to 10°C over about 2.5hr. The reaction was quenched by the addition of 100ml saturated NH₄Cl and 100 ml H₂O. Extracted once with EtOAc (300ml) and eight times with CH₂Cl₂ (150 ml). Dried over solid MgSO₄ and filtered. Concentrated and flashed over silica gel chromatography (3 to 10% MeOH (NH₃)/CH₂Cl₂) to obtain 85 (11g, yield: 38%).


Step 2

To the mixture of 85 (9.2 g) and MnO₂ (42 g) was added 200ml CH₂Cl₂, and the mixture was stirred at room temperature overnight. Additional MnO₂ (20g) was added and the reaction was stirred another 24hrs. The MnO₂ was filtered off and the reaction was concentrated and flashed over silica gel (5% and 10% MeOH (NH₃)/CH₂Cl₂) to give 86 (3.1g, yield: 33%).


Step 3

In a manner similar to that described in Example 7, step 2, 86 (3.1g) was converted to 87 (2.0 g, yield: 68%).


- 61 -

Step 4

In a manner similar to that described in Example 7, step 3, 4, 5, and 6, 87 was converted to 88.


Example 20**Step 1**

To the solution of compound 89 in CH_2Cl_2 (20ml) at 0°C was added m-CPBA (0.54g) and the reaction was stirred at 0°C for 25 min. and then at room temperature stirred for 2 hrs. 40% NH_4OH (12ml) was added and the mixture was stirred for 30min. Separated and extracted the aqueous layer with CH_2Cl_2 (10ml). Dried (MgSO_4), filtered and concentrated in vacuo. Flash chromatography (5% $\text{MeOH}(\text{NH}_3)/\text{CH}_2\text{Cl}_2$) gave 90 (0.67g, 80%).

Step 2


To the solution of 90 (0.65g) in CH_2Cl_2 (6 ml) at -10°C was added TFA (6ml) and the reaction was stirred for 1hr from -10°C to 0°C . Concentrated down and azeotroped twice with toluene (20ml), and concentrated to dryness to obtain 91 as a gummy oil which was used as is.

- 62 -

Step 3


In a manner similar to that described in Example 7, steps 5 and 6, **91** was converted to **92**.

5


Example 21**Step 1**

To a solution of **93** (5.17 g, 22.7 mmol) in THF (100 mL) at -50°C was added $s\text{-BuLi}$ (38.4 mL of a 1.3M solution in hexane, 49.9 mmol) dropwise. After 1.5h at -40°C , the reaction was recooled to -50°C and **95** (4.84 g, 22.7 mmol) in THF (20 mL) was added. After 2.75 h at -50°C , glacial acetic acid was added followed by saturated aqueous NH_4Cl . The mixture was warmed to room temperature and the layers were separated. The aqueous layer was extracted with EtOAc. The combined organic layers were dried (MgSO_4) filtered and concentrated to give a residue that was purified by flash column chromatography (1% to 3% MeOH/NH_3 in CH_2Cl_2) to give **95** (6.35 g, 63%).


- 63 -

Step 2

In a manner similar to that described in Example 12, step 3, **95** (5.34 g, 12.11 mmol) was converted to **96** (4.71 g, 75%).


Step 3

In a manner similar to that described in Example 6, step 4, **96** (3.7 g, 8.43 mmol) was converted to **97** (3.08 g, >100%) which was used as is in the next step.

Step 4


Compound **97** (0.7 g, 2.25 mmol), $\text{H}_2\text{NOCH}_3 \cdot \text{HCl}$ (0.94 g, 11.23 mmol) and NaOAc (1.47 g, 17.97 mmol) were combined in 1-pentanol (20 mL) and water (2 mL) and heated to reflux for 2 days. The reaction was cooled to room temperature and 0.5 N NaOH was added. The EtOH was removed in vacuo, additional water (15 mL) was added, and the reaction extracted with 10% EtOH in CH_2Cl_2 (180 mL total volume). The combined organic extracts were dried and concentrated to give **98** (0.55 g, 92%).

- 64 -

Step 5

In a manner similar to that described in Example 6, steps 5, 6, and 7, **98** was converted to **99**.


5

Example 22**Step 1**

prepared according to:
J. Org. Chem., 1968,
33(6), 2388


A solution of 2.2 g (9.5 mmol) of **100** in 75 mL of glacial acetic acid was
 10 hydrogenated in the presence of 0.5 g of 10% w/w platinum-on-charcoal for 5 h. The
 reaction mixture was filtered to remove the catalyst and the filtrate was concentrated
 by evaporation under reduced pressure to produce a solid residue which was basified
 with 0.5N NaOH and extracted with methylene chloride (CH_2Cl_2). Methylene chloride
 extracts were dried over anhydrous MgSO_4 and concentrated. The residue was
 15 purified by flash chromatography eluted with 10 - 30% of 7N NH_3 -MeOH in CH_2Cl_2 to
 give 0.82 g of **101** (mp 158-163 $^\circ\text{C}$). LCMS m/z 240 (MH^+).

- 65 -

Step 2


A mixture of 0.12 g (0.52 mmol) of **101**, 0.2 g (0.52 mmol) of **5A**, 0.67 g (0.5 mmol) of 1-hydroxybenzotriazole hydrate (HOBt), and 0.11 g (0.57 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (DEC) in 7 mL of anhydrous dimethylformamide (DMF) was stirred at ambient temperature for 18 h. The mixture was diluted with water and the resulting precipitate was filtered to produce 0.26 g of **102** as a white solid (mp 110-115 °C). LCMS m/z 557 (MH⁺).

- 66 -

Step 3

To a stirred solution of 0.34 g (2.7 mmol) of oxalyl chloride in 3 mL of anhydrous CH_2Cl_2 at -70°C was added 0.44 g (5.7 mmol) of anhydrous methylsulfoxide in 2 mL of CH_2Cl_2 . After being stirred at -70°C for 10 minutes, the reaction mixture was added 1.2 g (2.15 mmol) of **102** in 10 mL of CH_2Cl_2 . The stirred mixture was kept at -70°C for 0.5 h, mixed with 1.8 mL (13 mmol) of triethylamine, and then allowed to warm up to ambient temperature by itself. The mixture was diluted with water and extracted with CH_2Cl_2 . Organic extracts were washed with brine, dried over anhydrous MgSO_4 and concentrated to produce 1.18 g of **103** as a glass. LCMS m/z 555 (MH^+).

- 67 -

Step 4


A solution of 0.8 g (1.44 mmol) of **103** and 0.6 g (7.2 mmol) of methoxylamine hydrochloride in 40 mL of ethanol and 40 mL of pyridine was heated under reflux for 18 h. The mixture was concentrated and the residue was taken up in ethyl acetate/ether and washed with water. The organic solution was dried over anhydrous MgSO_4 and concentrated to 0.65 g of viscous residue which was dissolved in 8 mL of trifluoroacetic acid and 8 mL of CH_2Cl_2 and stirred at ambient temperature for 18 h. The solution was concentrated and the residue was basified with 1N NaHCO_3 and extracted with ethyl acetate. Organic extracts were washed with brine, dried over anhydrous MgSO_4 and concentrated to a gummy residue. Purification of this residue by flash chromatography with 5 - 8% of 7N NH_3 -MeOH in CH_2Cl_2 produced 0.151 g of **104** as a gum, LCMS m/z 484 (MH^+) and 0.146 g of **105** as a glass, LCMS m/z 556 (MH^+).

- 68 -

Mixing a solution of 0.056 g of the free base of **104** in ethyl acetate with a solution of 0.04 g of maleic acid in ethyl acetate produced a precipitate which was isolated by filtration to give 0.06 g of a dimaleate salt of **104** (mp 155-160 °C).

5 **Example 23**

Step 1




J. Med. Chem., 1976,
19, 360

2.4 g (10. mmol) of **106** were reduced in the similar manner as that described in Example 22, step 1 to give 1.5 g of **107** as a semi-solid. LCMS m/z 240 (MH⁺).

10


Step 2



1.5 g (6.31 mmol) of **107** were coupled with **3** in the similar manner as that described in Example 22, step 2 to give 3 g of **108** as a solid (mp 104-106 °C).

15 LCMS m/z 557 (MH⁺).


- 69 -

Step 3

1.17 g (2.1 mmol) of **108** were oxidized in the similar manner as that described in Example 22, step 3 to give 0.7g of **109** as a glass. LCMS m/z 557 (MH⁺).

- 70 -


Step 4



0.32 g (0.58 mmol) of **109** were reacted with 0.6 g (7.2 mmol) of methoxylamine hydrochloride in the same manner as that described in Example 22, step 4 to provide 0.065 g of **110** as a gum, LCMS m/z 484 (MH^+) and 0.12 g of **111** as a glass, LCMS m/z 556 (MH^+).

Example 24


Step 1



**J. Med. Chem., 1994,
37 (16), 2537**


- 71 -

A mixture of 18 g (74 mmol) of **112**, 7.2 g (74 mmol) of N,O-dimethylhydroxylamine hydrochloride, 19.4 g (15 mmol) of N,N-diisopropylethylamine, 1.1 g (8 mmol) of HOBt and 14.2 g (74 mmol) of DEC in 80 mL of anhydrous DMF was stirred at ambient temperature for 18 h. The mixture was diluted with water and extracted with ethyl acetate. Organic extracts were washed with 1% NaHCO₃ and brine, dried over anhydrous MgSO₄ and concentrated to give 15.5 g of **113** as an oil. LCMS m/z 287 (MH⁺).

Step 2


To a stirred solution of 2.9 g (18 mmol) of 2-bromopyridine in 30 mL of anhydrous THF at -78°C was added 7.5 mL of 2.5M solution of n-BuLi in hexane dropwise for 0.5 h. After being stirred at -78°C for 1 h, the reaction mixture was added a solution of 5.1 g (17.8 mmol) of **113** in 15 mL of THF. The mixture was allowed to stir at ambient temperature for 48 h, mixed with saturated aqueous NH₄Cl and extracted with ether. Organic extracts were washed with brine, dried over anhydrous MgSO₄ and concentrated to produce 5.7 g of **114** as an oil. LCMS m/z 305 (MH⁺).

20


Step 3

A solution of 3.15 g (10.4 mmol) of **114** and 3.47 g (41.6 mmol) of methoxylamine hydrochloride in 30 mL of ethanol and 30 mL of pyridine was heated under reflux for 18 h. The mixture was concentrated and the residue was taken up in ether and washed with water. The organic solution was dried over anhydrous MgSO₄ and concentrated to give 2.5 g of **115** as an oil. LCMS m/z 334 (MH⁺).

- 72 -


Step 4

A solution of 2.4 g (7.2 mmol) of **22** in 20 mL of CH_2Cl_2 and 20 mL of trifluoroacetic acid was stirred at ambient temperature for 1 h. The solution was concentrated. The residue was basified with saturated aqueous NaHCO_3 and extracted with CH_2Cl_2 . Organic extracts were washed with brine, dried over anhydrous MgSO_4 and concentrated to give 1.41 g of **23** as a glass. LCMS m/z 234 (MH^+).


10 **Step 5**

A mixture of 0.466 g (2 mmol) of **116**, 0.517 g (2.2 mmol) of **5A**, 0.276 g (2 mmol) of HOBt and 0.46 g (2.4 mmol) of DEC in 20 mL of anhydrous DMF was stirred at ambient temperature for 18 h. The mixture was concentrated by evaporation under reduced pressure at bath temperature of 25-45°C and the residue was chromatographed with 4% (7N $\text{NH}_3/\text{CH}_3\text{OH}$) in CH_2Cl_2 to produce 0.48 g of syrup which was dissolved in 15 mL of EtAc-EtOH (3:1 v) and mixed with a solution of 0.26 g of maleic acid in 10 mL of EtAc-EtOH (1:1). The resulting precipitate was filtered to produce 0.35 g of the maleate salt of **117** (mp 160-163 °C). LCMS m/z 451 (MH^+).

- 73 -

Example 25**Step 1**


To a stirred solution of 4.16 g (20 mmol) of 1A in 80 mL of anhydrous THF at
 5 -78°C was added dropwise 17 mL of 2.5M solution of n-BuLi in hexane for 25
 minutes. After being stirred from -78°C to room temperature for 1 h, the reaction
 mixture was added a solution of 6 g (22 mmol) of 26 in 100 mL of anhydrous THF and
 kept at room temperature for 18 h. The mixture was mixed with saturated aqueous
 NH₄Cl and extracted with EtAc. Organic extracts were washed with brine, dried over
 10 anhydrous MgSO₄ and concentrated to produce 6.1 g of 118 (mp 146-149 °C). LCMS
 m/z 420 (MH⁺).

Step 2


15 A solution of 3.71 g (8.8 mmol) of 118 and 3.7 g (44 mmol) of methoxyamine
 hydrochloride in 40 mL of pyridine and 40 mL of ethanol was heated under reflux for 2
 days. The mixture was concentrated and the residue was taken up in CH₂Cl₂ and
 washed with saturated aqueous NaCl. Organic solution was dried over anhydrous
 MgSO₄ and concentrated to give 2.6 g of 119 as a glass. LCMS m/z 421 (MH⁺).

20


- 74 -

Step 3

A solution of 0.9 g (2.14 mmol) of **119** in 10 mL of CH_2Cl_2 and 10 mL of trifluoroacetic acid was stirred at ambient temperature for 2 h. The solution was concentrated. The residue was taken up in CH_2Cl_2 , washed with saturated NaHCO_3 and brine, dried over anhydrous MgSO_4 and concentrated to a solid residue which was triturated with CH_3CN and filtered to produce 0.29 g of **120** (mp 200-205 °C). LCMS m/z 321 (MH^+).


Step 4

0.1 g (0.31 mmol) of **120** and 0.83 g (0.35) of **5A** were coupled in the same manner as that described in Example 24, step 5 to produce 0.12 g of the maleate salt of **121** (mp 170-173 °C). LCMS m/z 538 (MH^+).



Example 26**Step 1**

In a similar manner to that described in Example 6, step 7, **122** (0.26 g, 0.41 mmol) was converted to **123** (0.08 g, 40%).

- 75 -


Example 27**Step 1**

To a suspension of LAH (0.83 g, 22 mmol) in ether (20 mL) at 0° C was added
5 **124** (3.2 g, 17.5 mmol) in THF (15 mL) dropwise. The reaction was stirred at 0° C for
1.5 h, and quenched by the addition of water (0.8 mL), 20% aqueous NaOH (0.8 mL),
and water (2.4 mL). The mixture was stirred for 15 min and filtered and the filter cake
washed with CH₂Cl₂. The filtrate was concentrated to give an oil which was
dissolved in ether (30 mL) and washed with brine and dried (MgSO₄). Filtration and
10 concentration in vacuo gave **125** (2.5 g) which was used without further purification.

Step 2**Step 3**


In a similar manner to that described in Example 22, step 3 and Example 1,
steps 4, 5, and 6, **125** was converted to **126**.

- 76 -

Step 4

In a similar manner to that described in Example 6, step 5, 126 was converted to 127.











5











Step 5













10 In a similar manner to that described in Example 6, step 7, 127 was converted to 128.













The compounds in Table 1 (first column) are prepared from the compounds in the last column of Table 1 by following essentially the same procedures as in the examples described above. In Table 1 "Cmpd. No." stands for "Compound Number".













TABLE 1













Cmpd. No.	STRUCTURE	Mass Spec. $[M^+H]^+$	Starting Material
200		470.1	 •HCl
201		456.1	 •HCl
202		456.1	 •HCl
203		531.1	 •HCl
204		499.1	 •HCl













205		497.1	 •HCl
206		517.1	 •HCl
207		549.1	 •HCl
208		599.1	 •HCl
209		568.1	 •HCl













210		565.1	 ·HCl
211		483	 ·HCl
212		484.1	 ·HCl
213		583.1	 ·HCl
214		552.1	 ·HCl
215		471	 ·HCl













216		512	
217		512	
218		504	
219		454	
220		470	
221		456	













222		456	
223		495	
224		470	
225		470	
226		504	
227		484	















228		472	
229		486	
230		572	
231		505	
232		452	
233		518	

















234		450	
235		442	
236		423	
237		423	
238		436	
239		451	











240		423	
241		423	
244		435	
245		519	
246		451	
247		421	

248		438	
249		452	
250		487	
251		543	
252		501	
253		457	

254		471	
255		465	
256		465	
257		422	
258		406	
259		455	



260		484	
261		443	
262		440	
263		441	
264		427	
265		427	
266		518	

267		490	 •HCl
268		455	 •HCl
269		439	 •HCl
270		407	 •HCl
271		421	 •HCl
272		407	 •HCl
273		455	 •HCl
275		425	 •HCl

278		425	 •HCl
279		439	 •HCl
280		470	 •HCl
281		469	 •HCl
282		504	

- 90 -


The isomers 246A and 253A, below, can be separated from 246 and 253, respectively, above, by techniques well known to those skilled in the art.



5

Example 28

Step 1




10 To a solution of 1.00 g (8.13 mmol) of pyrimidine aldehyde 67 (Step 2 of Example 13) in 40 ml of CH_2Cl_2 was added 1.36 mL (10.58 mmol) of ethyl isonipecotate and 2 drops of acetic acid. The mixture was stirred for 40 min. at room temperature, after which 2.58 g (12.17 mmol) of $\text{NaBH}(\text{OAc})_3$ was added. The reaction mixture was then stirred for 20 h at room temperature, diluted with aqueous

15 NaOH (pH adjusted to 11) and extracted with CH_2Cl_2 . Organic phase was dried and concentrated, and the residue was flash chromatographed (4-8% ca. 3 N NH_3 in

- 91 -


MeOH/ CH₂Cl₂) to produce 1.55 g (5.87 mmol; 72%) of amine **285** as a yellowish solid.

Step 2

**286**

To a solution of 3.83 g (14.51 mmol) of ester **285** in 60 ml of 3 : 1 : 1 mixture of THF – MeOH – H₂O was added 1.22 g (29.02 mmol) of LiOH monohydrate. The reaction mixture was stirred at room temperature overnight, concentrated, and the residue was dried under high vacuum to produce 3.84 g of crude acid **286** lithium salt as a yellow solid. Material could be used directly or could be purified by passing through a silica gel plug eluting with ca. 3 N NH₃ in MeOH.

Step 3

**287**

To a mixture of 3.32 g (14.05 mmol) of acid **286** and 4.07 g (14.05 mmol) of 4-[(E)-(methoxyimino)-2-pyridinylmethyl]piperidine dihydrochloride (see Compound **447** below) in 40 mL of DMF was added 8.94 mL (70.25 mmol) of 4-ethylmorpholine and 14.0 mL (23.52 mmol) of 50 wt. % solution of 1-propanephosphonic acid cyclic anhydride in ethyl acetate. The reaction mixture was stirred for 4.5 h at 50°C followed by 14 h at room temperature. Concentration of the mixture was followed by exposure to high vacuum for 24 h to remove remaining DMF. The residue was partitioned between aqueous NaOH and CH₂Cl₂, organic phase was separated, dried and


- 92 -

concentrated, and the residue was flash chromatographed (5-15% ca. 3 N NH₃ in MeOH/ CH₂Cl₂) to produce 4.60 g (10.51 mmol; 75 %) of amide **287** as a light tan foam. MS 438 (M+1).

5 **Example 29**


Step 1

Reference: *J. Heterocyclic Chem.*, **1966**, 3, 252.



3,4 Pyridine-dicarboximide **288** (10.0 g; 67.5 mmoles) was dissolved in 162 g.
10 of 10% aqueous NaOH and the solution was cooled to an internal temperature of 7 °C
in an ice-salt bath. Bromine (3.6 ml; 70 mmoles) was added dropwise. After the
addition, the solution was heated for 45 minutes at a bath temperature of 80-85 °C.
The yellow solution was then cooled to an internal temperature of 37 °C, then 17 ml of
glacial acetic acid were added dropwise to a pH of 5.5. The resulting mixture was
15 saved overnight in a refrigerator. The solid formed was filtered and washed with 5 ml
of water and 5 ml of methanol. The reaction yielded 6.35 g. of product **289** melting at
280-285 °C (decomp.).


Step 2




20 Solid Compound **289** (9.5 gr.; 69 mmoles) was carefully added in three aliquots
to a slurry of lithium aluminum hydride (9.5 gr.; 250 mmoles) in 200 ml of dry
tetrahydrofuran. The resulting hot mixture was stirred at room temperature for two
days. After cooling in an ice bath, the reaction was quenched with very careful
25 sequential dropwise addition of 10 ml of water, followed by 10 ml of 15% aqueous

- 93 -

NaOH, then by 30 ml of water. The resulting solid was filtered through a pad of Celite and washed several times with THF. The oil obtained after evaporation of the solvent, solidified on standing. The reaction mixture was purified by flash chromatography on silica gel using 5%MeOH(NH₃)/EtOAc as eluent yielding 6.21
5 (72%) of Compound 290 . LC-MS: m/z = 125 (M+1).

Step 3


Manganese dioxide (29 gr.; 334 mmoles) was added, in one portion, at room
10 temperature, to a suspension of 3-amino-4-hydroxymethyl pyridine 290 (5.0 gr.; 40.3 mmoles) in 500 ml of chloroform with good stirring. After two days, the solid is filtered through a pad of Celite and washed with chloroform. Removal of the solvent using reduced pressure yielded 4.2 grams (85%) of Compound 291 as a yellow solid.

Step 4


A dry dichloromethane (400 ml) solution of ethyl isonipecotate (12.5 gr.; 79.5 mmoles) and 3-amino pyridine 4-carboxyaldehyde 291 (3.33 gr.; 27.3 mmoles) was stirred at room temperature for one hour, then 60 grams of activated 3Å molecular
20 sieves were added. The mixture was stirred for an additional 90 minutes, then 20 grams (96.4 mmoles) of sodium triacetoxo borohydride was added at room temperature in one portion. After stirring for three days, the solid was filtered through a pad of Celite and washed with dichloromethane. The solution was stirred for 15

- 94 -

minutes with 100 ml of saturated aqueous sodium bicarbonate then separated from the aqueous layer. The organic layer was washed two more times with saturated aqueous sodium bicarbonate, then with brine and dried with anhydrous sodium sulfate. After evaporation of the solvent, the resulting oil was purified by flash chromatography on silica gel using EtOAc:Hexanes:MeOH(NH₃) as eluent. The procedure yielded 6.8 gr.(94%) of Compound 292 . FAB-MS: m/z = 264 (M+1).

Step 5


10 Ethyl 1-[(3-amino-4-pyridinyl)methyl]-4-piperidinecarboxylate 292 (4.75 gr.; 18.04 mmoles) was stirred for 24 hours at room temperature with 1.51 gr. (36 mmoles) of lithium hydroxide monohydrate in 75 ml of methanol. Removal of the solvent using reduced pressure yielded Compound 293 as a white solid.

15 **Step 6**

20 4-(2-pyridinylcarbonyl)piperidine 28 (Step 4 in Example 6) (0.3 gr.; 1.58 mmoles), lithium 1-[(3-amino-4-pyridinyl)methyl]-4-piperidinecarboxylate 293 (0.34 gr.; 1.4 mmoles), DEC (0.38 gr.; 2.0 mmoles), and HOBT (0.27 gr.; 2.0 mmoles) were stirred at room temperature in 10 ml of dry DMF for two days. The reaction was

- 95 -

quenched with 50 ml. of 0.5 N aqueous NaOH, then the solution was extracted with dichloromethane. The combined extracts were washed with brine and dried over anhydrous sodium sulfate. The product **295** was isolated by flash chromatography on silica gel using EtOAc:Hexanes:MeOH(NH₃) (50:45:5) as eluent. Yields: 0.27 gr. (47%). FAB-MS: m/z = 408 (M+1).

Step 7

1-[[[1-[(3-amino-4-pyridinyl)methyl]-4-piperidiny]carbonyl]-4-(2-pyridinylcarbonyl)piperidine **295** (0.196 gr.; 0.48) and methoxyamine hydrochloride (0.401 gr. 4.8; mmols) were heated, under N₂, at a bath temperature of 70°C for 24 hours in 6.0 ml of dry pyridine. After removing the pyridine using reduced pressure, the residue was treated with saturated aqueous sodium bicarbonate. The resulting mixture was extracted several times with dichloromethane. The combined extracts were washed with brine and dried over anhydrous sodium sulfate. The reaction mixture was purified by silica gel preparative thin layer chromatography. The plates were eluted with EtOAc:Hexanes:MeOH(NH₃) (60:35:5) and the product **296** was extracted with 10% MeOH(NH₃)/EtOAc. Yields: 0.15 gr. (71%). FAB-MS: m/z = 437 (M+1).


Example 30 Step 1

A mixture of **297** (1 g, 10 mmol) in 1:1 water-dioxane (50 mL) was treated with Et₃N (4 mL, 13 mmol) and BOC₂O (2.8 g, 13 mmol) at 4°C and allowed to warm to 20°C for one day. The solvent was then removed in vacuo. The residue was taken up in 1:1 water-ethyl acetate and the organic layer was discarded. The aqueous

- 96 -


layer was acidified with 1 N aqueous HCl and extracted three times with ethyl acetate. The combined organic phases were washed with water and brine, dried (Na₂SO₄), and concentrated to give **298** as a white solid (1.8 g, 90%).

5 **Step 2**



A mixture of **298** (1.8 g, 9 mmol), N,O-dimethylhydroxylamine hydrochloride (2.6 g, 27 mmol), EDCI (5 g, 27 mmol), HOBt (0.1 g, 1 mmol), and DIPEA (12.5 mL, 72 mmol) in DMF (30 mL) was stirred at 20°C overnight. The reaction was then
10 concentrated to half volume in vacuo, poured onto water, and extracted three times with ethyl acetate. The combined organic phases were washed with saturated aqueous NH₄Cl, saturated aqueous NaHCO₃, water and brine, dried (Na₂SO₄), and concentrated to give **299** as a clear oil (2.1 g, 98%).

15 **Step 3**




To a solution of 2-bromopyridine (1.2 mL, 12 mmol) in THF (60 mL) at -78°C was added n-BuLi (8 mL of a 1.6 M solution in hexanes, 12 mmol) dropwise over 15 min. After stirring for an additional 30 min at -78 °C, a solution of **299** (1 g, 4 mmol) in
20 THF (20 mL) was slowly added. The reaction was then heated to 60 °C for 1 h. After cooling to 20 °C, the reaction was diluted with ether, quenched with saturated aqueous Na₂SO₄, and dried with solid Na₂SO₄. The mixture was filtered through a plug of solid Na₂SO₄ and concentrated in vacuo. Flash column chromatography (0–20% ethyl acetate–hexanes) yielded **300** as a yellow oil (0.12 g, 11%).

25

- 97 -






Step 4

















Following procedures similar to those of Steps 4 to 7 of Example 6, compound 301 was obtained. MS 409 (M+1).








5








Following procedures similar to those described in the examples above, the compounds in Table 2 were prepared.







TABLE 2		
Compound	STRUCTURE	MS (M+1)
302		430
303		421
304		505
305		505
306		471






307		426
308		408
309		442
310		437
311		437
312		458
313		402








314		487
315		438
316		467
317		424
318		451
319		430
320		523








321		453
322		453
323		410
324		413
325		439
326		466
327		453








328		453
329		424
330		453
331		438
332		488
333		437
334		437






335		479
336		452
337		466
338		438
339		465
340		465






341		513
342		452
343		550
344		499
345		451




346		451
347		451
348		451
349		452
350		455
351		455
352		422

353		422
354		492
355		438
356		437
357		424
358		510
359		539

360		453
361		409
362		438
363		426
364		422
365		483
366		483

367		497
368		465
369		479
370		479
371		493















372		564
373		517
374		568
375		426
376		455















377		456
378		452
379		427















- 111 -















If one were to follow procedures similar to those described in the examples above, the compounds in the "Structure" column of Table 3 would be obtained using the starting material listed in Table 3. Each compound in Table 3 is a mixture of oxime isomers, as represented by the \sim bond between the oxime nitrogen and the
5 OH or OCH₃ moiety. In Table 3 "CMPD" stands for "Compound".

TABLE 3















CMPD	Structure	Starting Material
380		 •HCl
381		 •HCl
382		 •HCl
383		 •HCl
384		
385		 •HCl
386		















CMPD	Structure	Starting Material
387		
388		
389		
390		
391		
392		
393		















CMPD	Structure	Starting Material
394		
395		
396		
397		
398		
399		
400		









CMPD	Structure	Starting Material
401		
402		
403		
404		
405		
406		
407		

CMPD	Structure	Starting Material
408		
409		
410		
411		
412		
413		
414		


CMPD	Structure	Starting Material
415		
416		
417		
418		
419		
420		
421		

CMPD	Structure	Starting Material
422		
423		
424		
425		
426		
427		
428		


CMPD	Structure	Starting Material
429		
430		
431		
432		
433		
434		
435		

CMPD	Structure	Starting Material
436		
437		
438		
439		


- 121 -

Example 31**Step 1**

- 5 To a solution of LDA (233 mL, 2.0 M in THF/heptane/ethylbenzene, 0.466 mol) in THF (300mL) at 0 °C was added, dropwise over 1.0 h, a solution of compound 440 (100g, 0.389 mol) in THF (~ 400 mL). The red-orange solution was stirred at 0 °C for 30 min, and then transferred by cannula to a pre-cooled (0 °C) solution of N-fluorobenzenesulfonimide (153 g, 0.485 mol) in dry THF (~ 600 mL). The reaction
- 10 mixture was stirred at 0 °C for 30 min, and then at rt for 18 h. The total solvent volume was reduced to approximately one third, and EtOAc (~ 1L) was added. The solution was washed successively with water, 0.1 N aq. HCl, saturated aq. NaHCO₃, and brine. The organic layer was dried over MgSO₄, filtered, and concentrated under reduced pressure to yield a crude liquid. Separation by flash chromatography (6:1
- 15 hexanes-EtOAc) gave compound 441 (93.5 g, 87%).

Step 2


- 20 In a manner similar to that described in Example 6, Step 4, compound 441 was converted to compound 442.

Step 3

25


- 122 -

In a manner similar to that described in Example 1, Step 4, compound **442** was converted to compound **443**.

Step 4


In a manner similar to that described in Example 1, Step 5, compound **443** was converted to compound **444**.

10

Step 5

In a manner similar to that described in Example 6, Step 5, compound **5** was converted to compound **445**.


15

Step 6


20 In a manner similar to that described in Example 6, Step 6, compound **445** was converted to compound **446**.

- 123 -


In the above examples, the compound 4-[(E)-(methoxyimino)-2-pyridinylmethyl]piperidine dihydrochloride:



- 5 can be used to prepare the compounds of this invention, for example, see Examples 6 and 28. Preferably, Compound 447 is prepared from a compound of formula:



and from a compound of Formula 449:




10

R^{50} is an alkyl or aryl group, f is 0 to 4, R^{51} is an alkyl group, and Q is a halo group, wherein said alkyl, aryl, and halo groups are as defined above.


Compound 447 can be prepared from 448 and 449 by:

- 15 (a) converting the compound of formula 449 into its Grignard form (449A):



- 20 (b) reacting the compound of formula 448 with the compound of formula 449A to obtain a compound of formula 450:


- 124 -



(c) reacting the compound of formula 450 with a suitable alkyl chloroformate of formula 451




to yield a compound of formula 452:




10

(d) forming the salt (formula 453):



15 (e) reacting the compound of formula 453 with an alkoxyamine ($\text{NH}_2\text{OR}^{51}$) or its hydrochloride to form an oxime of formula 454:




(f) isomerizing the compound of formula 454 by treatment with a strong acid and simultaneously converting to the desired acid salt of Formula 454 with an enriched E isomer, wherein the E isomer predominates over the Z-isomer by at

20

- 125 -


least a 90:10 ratio. When $f=0$, R^{51} is methyl, and the acid used for isomerization is HCl in the compound of formula 454, the final product is the compound of formula 447.

This preparation can be represented as follows:



- 126 -

Following the above process the Compound 447 can be prepared as follows:



The conversion of compound 461 to 447 predominantly yields the E-isomer of compound 447 in high stereochemical purity and high yields. Isomerization of a mixture of phenyl compounds by acid catalysis is discussed by T. Zsuzsanna *et al*, *Hung.Magy.Km.Foly.*, 74(3) (1968), 116-119.

The above process starts with Compound 449. In step 1, a 4-halo-1-alkylpiperidine (or a 4-halo-1-arylpiperidine) is converted to its Grignard analog (449A) by reacting with magnesium. The reaction is performed generally at temperatures of about -10°C to reflux. Generally a hydrocarbon solvent such as, for example, toluene, xylene, chlorobenzene, dichlorobenzene and the like, or mixture of hydrocarbons listed above with an ether, such as, for example, a $\text{C}_5\text{-C}_{12}$ alkyl ether, 1,2-dimethoxyethane, 1,2-diethoxyethane, diglyme, 1,4-dioxane, tetrahydrofuran and the like are suitable for this reaction. The solution is cooled to around -10°C to about 10°C and then reacted with a suitable 2-cyanopyridine (448), for about 10-120 minutes. Examples of suitable 2-cyanopyridines are 2-cyanopyridine, 4-methyl-2-

- 127 -

cyanopyridine, 4-ethyl-2-cyanopyridine, 4-phenyl-2-cyanopyridine, and the like. Preferred are 2-cyanopyridine and 4-methyl-2-cyanopyridine. The Grignard compound is used generally in about 1-4 molar equivalents with respect to the compound of formula 448, preferably in about 1-3 molar equivalents and typically in about 1.5-2.5 molar equivalents. The product of formula 450 may be isolated by procedures well known in the art, such as, for example, treatment with an acid (e.g. HCl); preferably in a suitable solvent (e.g., tetrahydrofuran or ethyl acetate).

The product of Formula 450 may then be reacted with an alkyl chloroformate in the next step. Suitable alkyl chloroformates are, for example, methyl chloroformate, ethyl chloroformate, propyl chloroformate, and the like, with the preferred being methyl chloroformate or ethyl chloroformate. Generally a hydrocarbon solvent such as, for example, toluene, xylene, chlorobenzene, dichlorobenzene and the like, or mixture of a hydrocarbons listed above with an ether such as, for example, a C₅-C₁₂ alkyl ether, 1,2-dimethoxyethane, 1,2-diethoxyethane, diglyme, 1,4-dioxane, tetrahydrofuran and the like is suitable for this reaction. The reaction is generally performed at about 25-100°C, preferably about 40-90°C and typically about 50-80°C, for about 1-5 hours. After the reaction, generally the generated acid is washed off and the product of formula 452 may be isolated by organic solvent extraction.

The compound of Formula 452 may then be converted into its acid salt by treatment with an acid such as, for example, sulfuric acid, hydrochloric acid, trifluoroacetic acid and the like, generally in a solvent at temperatures between ambient and reflux of the solvent. Suitable solvents include hydrocarbons such as, for example, toluene, xylene, chlorobenzene, dichlorobenzene and the like. There being two nitrogen atoms in the compound of Formula 452, the salt generally has 2 moles of acid to a mole of compound 452.

The compound of formula 453 may then be converted to an alkyloxime of formula 454 by reacting it with an alkoxyamine (or its hydrochloride), usually in aqueous solution form. Suitable alkoxyamines are, for example, methoxyamine, ethoxyamine and the like. Methoxyamine is preferred. The alkoxyamine (or its hydrochloride) is employed generally in about 1 to about 4 molar equivalents, preferably in about 1 to about 3 molar equivalents, and typically in about 1 to about 2 molar equivalents. Generally, the reaction is catalyzed by a weak acid such as, for example, acetic acid, formic acid and the like, or mixtures thereof. A cosolvent such

- 128 -

as, for example, methanol, ethanol, isopropanol, n-butanol and the like, or mixtures thereof may be added. The product of formula 454, after work-up, is a mixture of the Z- and the E-isomers, whose ratio may be analyzed for its stereochemical make-up, using techniques well known in the art such as, for example, HPLC.

5 Treating the compound of formula 454 with a strong acid under the reaction conditions described below isomerizes the mixture of the Z and the E-isomers into predominantly the E-isomer. Generally, the compound of formula 454 may be dissolved in a solvent such as, for example, ethanol, methanol, isopropanol, n-butanol and the like, ether such as methyl tert-butyl ether, tetrahydrofuran and the like,
10 hydrocarbon such as, for example, heptane, hexane, toluene and the like, nitrile such as, for example, acetonitrile, benzonitrile and the like, or mixtures of such solvents. The dissolved compound is then treated with a strong acid such as, for example, HCl, HBr, H₂SO₄ and the like, at temperatures in the range of 20 to 100°C for about 1-20 hours. The acid is employed generally in about 1 to about 8 molar equivalents,
15 preferably in about 1 to about 6 molar equivalents, and typically in about 2 to about 4 molar equivalents. Work-up typically forms predominantly the acid salt of the E-isomer of the compound of formula 454, which is in fact the compound of formula 447 when R⁵¹ = methyl, n=0 and the acid salt is HCl in 454.

The products of the various steps in the process described above may be
20 isolated and purified by conventional techniques such as, for example, filtration, recrystallization, solvent extraction, distillation, precipitation, sublimation and the like, as is well known to those skilled in the art. The products may be analyzed and/or checked for purity by conventional methods such as, for example, thin layer chromatography, NMR, HPLC, melting point, mass spectral analysis, elemental
25 analysis and the like, well known to those skilled in the art.

H₃-Receptor Binding Assay

The source of the H₃ receptors in this experiment was guinea pig brain. The animals weighed 400-600 g. The brain tissue was homogenized with a solution of 50
30 mM Tris, pH 7.5. The final concentration of tissue in the homogenization buffer was 10% w/v. The homogenates were centrifuged at 1,000 x g for 10 min. in order to remove clumps of tissue and debris. The resulting supernatants were then centrifuged at 50,000 x g for 20 min. in order to sediment the membranes, which were

- 129 -

next washed three times in homogenization buffer (50,000 x g for 20 min. each). The membranes were frozen and stored at -70°C until needed.

All compounds to be tested were dissolved in DMSO and then diluted into the binding buffer (50 mM Tris, pH 7.5) such that the final concentration was 2µg/ml with 0.1% DMSO. Membranes were then added (400 µg of protein) to the reaction tubes. The reaction was started by the addition of 3 nM [³H]R-α-methyl histamine (8.8 Ci/mmol) or 3 nM [³H]N^α-methyl histamine (80 Ci/mmol) and continued under incubation at 30°C for 30 min. Bound ligand was separated from unbound ligand by filtration, and the amount of radioactive ligand bound to the membranes was quantitated by liquid scintillation spectrometry. All incubations were performed in duplicate and the standard error was always less than 10%. Compounds that inhibited more than 70% of the specific binding of radioactive ligand to the receptor were serially diluted to determine a K_i (nM).

Compounds 23, 30, 31, 32, 33, 44, 45, 49, 50, 53, 54, 55, 56, 57A, 59, 75, 76, 83, 88, 92, 99, 104, 110, 117, 128, 200, 201, 203-215, 217-241, 244-246, 246A, 247-253, 253A, 254-273, 275, 278, 280-282, 287, 296, 301-310, and 312-379 had a K_i within the range of about 0.25 to about 370nM.

Preferred Compounds 23, 30, 31, 32, 33, 50, 53, 54, 55, 56, 57A, 59, 92, 212, 215, 218, 219, 220, 224, 225, 226, 227, 229, 233, 235, 237, 238, 246, 246A, 247, 248, 251, 253, 253A, 268-273, 275, 278-281, 287, 296, 301, 304-307, 309, 312, 314-318, 320-356, and 358-376 had a K_i within the range of about 0.25 to about 33nM.

Most preferred Compounds 30, 31, 32, 33, 54, 55, 56, 56A, 225, 237, 246A, 253A, 273, 280, 287, 296, 301, 304-307, 309, 312, 314-318, 320-348, 350-356, 359-372, and 374-376 had a K_i within the range of about 0.25 to about 16nM.

More preferred compound 32 had a K_i of 0.83nM.

More preferred compounds 54, 55, 253A, 287, 320 had a K_i within the range of about 1.05 to about 9.75nM.

For preparing pharmaceutical compositions from the compounds described by this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets and suppositories. The powders and tablets may be comprised of from about 5 to about 95 percent active ingredient. Suitable solid carriers are known in the art,

- 130 -

e.g. magnesium carbonate, magnesium stearate, talc, sugar or lactose. Tablets, powders, cachets and capsules can be used as solid dosage forms suitable for oral administration. Examples of pharmaceutically acceptable carriers and methods of manufacture for various compositions may be found in A. Gennaro (ed.), Remington's
5 Pharmaceutical Sciences, 18th Edition, (1990), Mack Publishing Co., Easton, PA.

Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injection or addition of sweeteners and opacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal
10 administration.

Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier, such as an inert compressed gas, e.g. nitrogen.

Also included are solid form preparations which are intended to be converted,
15 shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

The compounds of the invention may also be deliverable transdermally. The transdermal compositions can take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type
20 as are conventional in the art for this purpose.

Preferably the compound is administered orally.

Preferably, the pharmaceutical preparation is in a unit dosage form. In such form, the preparation is subdivided into suitably sized unit doses containing appropriate quantities of the active component, e.g., an effective amount to achieve
25 the desired purpose.

The quantity of active compound in a unit dose of preparation may be varied or adjusted from about 1 mg to about 150 mg, preferably from about 1 mg to about 75 mg, more preferably from about 1 mg to about 50 mg, according to the particular application.

30 The actual dosage employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage regimen for a particular situation is within the skill of the art. For

- 131 -

convenience, the total daily dosage may be divided and administered in portions during the day as required.

5 The amount and frequency of administration of the compounds of the invention and/or the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated. A typical recommended daily dosage regimen for oral administration can range from about 1 mg/day to about 300 mg/day, preferably 1 mg/day to 75 mg/day, in two to four divided doses.

10

The methods of this invention described above using a compound of Formula I also include the use of one or more compounds of Formula I, and the methods of this invention described above using a compound of Formula I in combination with an H₁ receptor antagonist also include the use of one or more compounds of Formula I in combination with one or more H₁ receptor antagonists.


15

While the present has been described in conjunction with the specific embodiments set forth above, many alternatives, modifications and variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.

20

WHAT IS CLAIMED IS:

1. A compound of the formula:



or a pharmaceutically acceptable salt or solvate thereof, wherein:

- (1) R^1 is selected from:

- (a) aryl;
(b) heteroaryl;
(c) heterocycloalkyl;
(d) alkyl;
(e) cycloalkyl; or
(f) alkylaryl;



wherein said R¹ groups are optionally substituted with 1 to 4 substituents

independently selected from:

- (1) halogen;
- (2) hydroxyl;
- (3) lower alkoxy;
- (4) $-\text{CF}_3$;
- (5) $\text{CF}_3\text{O}-$;
- (6) $-\text{NR}^4\text{R}^5$;
- (7) phenyl;
- (8) $-\text{NO}_2$;
- (9) $-\text{CO}_2\text{R}^4$;
- (10) $-\text{CON}(\text{R}^4)_2$ wherein each R^4 is the same or different;
- (11) $-\text{S}(\text{O})_m\text{N}(\text{R}^{20})_2$ wherein each R^{20} is the same or different H or alkyl group, preferably C_1 to C_4 alkyl, most preferably C_1 - C_2 alkyl, and more preferably methyl;
- (12) $-\text{CN}$; or

- 133 -

(13) alkyl; or

(2) R^1 and X taken together form a group selected from:(3) X is selected from: $=C(O)$, $=C(NOR^3)$, $=C(NNR^4R^5)$,(4) M^1 is carbon;(5) M^2 is selected from C or N;(6) M^3 and M^4 are independently selected from C or N;

(7) Y is selected from: is $-CH_2-$, $=C(O)$, $=C(NOR^{20})$ (wherein R^{20} is as defined above), or $=C(S)$;

(8) Z is a $C_1 - C_6$ alkyl group;

(9) R^2 is a five or six-membered heteroaryl ring, said six-membered heteroaryl ring comprising 1 or 2 nitrogen atoms with the remaining ring atoms being carbon, and said five-membered heteroaryl ring containing 1 or 2 heteroatoms selected from: nitrogen, oxygen, or sulfur with the remaining ring atoms being carbon; said five or six membered heteroaryl rings being optionally substituted with 1 to 3 substituents independently selected from: halogen, hydroxyl, lower alkyl, lower alkoxy, $-CF_3$, CF_3O- , $-NR^4R^5$, phenyl, $-NO_2$, $-CO_2R^4$, $-CON(R^4)_2$ wherein each R^4 is the same or different, $-CH_2NR^4R^5$, $-(N)C(NR^4R^5)_2$, or $-CN$;

(10) R^3 is selected from:

(a) hydrogen;

(b) $C_1 - C_6$ alkyl;

(c) aryl;

(d) heteroaryl;


(e) heterocycloalkyl;

(f) arylalkyl;

(g) $-(CH_2)_e-C(O)N(R^4)_2$ wherein each R^4 is the same or different,(h) $-(CH_2)_e-C(O)OR^4$;

- 134 -

(i) $-(\text{CH}_2)_e-\text{C}(\text{O})\text{R}^{30}$ wherein R^{30} is a heterocycloalkyl group, such as, for example, morpholinyl, piperidinyl, piperazinyl or pyrrolidinyl, including



(j) $-\text{CF}_3$; or

5 (k) $-\text{CH}_2\text{CF}_3$;

wherein said aryl, heteroaryl, heterocycloalkyl, and the aryl portion of said arylalkyl are optionally substituted with 1 to 3 substituents selected from: halogen, $-\text{OH}$, $-\text{OCF}_3$, $-\text{CF}_3$, $-\text{CN}$, $-\text{N}(\text{R}^{45})_2$, $-\text{CO}_2\text{R}^{45}$, or $-\text{C}(\text{O})\text{N}(\text{R}^{45})_2$, wherein each R^{45} is independently selected from: H, alkyl, alkylaryl, or aryl wherein said aryl moiety is substituted with 1 to 3 substituents independently selected from $-\text{CF}_3$, $-\text{OH}$, halogen, alkyl, $-\text{NO}_2$, or $-\text{CN}$;

(11) R^4 is selected from: hydrogen, $\text{C}_1 - \text{C}_6$ alkyl, aryl, alkylaryl, said aryl and alkylaryl groups being optionally substituted with 1 to 3 substituents selected from: halogen, $-\text{CF}_3$, $-\text{OCF}_3$, $-\text{OH}$, $-\text{N}(\text{R}^{45})_2$, $-\text{CO}_2\text{R}^{45}$, $-\text{C}(\text{O})\text{N}(\text{R}^{45})_2$, or $-\text{CN}$; wherein R^{45} is as defined above;

(12) R^5 is selected from: hydrogen, $\text{C}_1 - \text{C}_6$ alkyl, $-\text{C}(\text{O})\text{R}^4$, $-\text{C}(\text{O})_2\text{R}^4$, or $-\text{C}(\text{O})\text{N}(\text{R}^4)_2$ wherein each R^4 is independently selected, and R^4 is as defined above;

(13) or R^4 and R^5 taken together with the nitrogen atom to which they are bound forms a five or six membered heterocycloalkyl ring;

20 (14) R^6 is selected from: alkyl, aryl, alkylaryl, halogen, hydroxyl, lower alkoxy, $-\text{CF}_3$, $\text{CF}_3\text{O}-$, $-\text{NR}^4\text{R}^5$, phenyl, $-\text{NO}_2$, $-\text{CO}_2\text{R}^4$, $-\text{CON}(\text{R}^4)_2$ wherein each R^4 is the same or different, or $-\text{CN}$;

(15) R^{12} is selected from: alkyl, hydroxyl, alkoxy, or fluoro;

(16) R^{13} is selected from: alkyl, hydroxyl, alkoxy, or fluoro;

25 (17) a is 0 to 2;

(18) b is 0 to 2;

(19) c is 0 to 2;

(20) e is 0 to 5;

(21) m is 1 or 2;


30 (22) n is 1, 2 or 3; and

- 135 -

(23) p is 1, 2 or 3, with the proviso that when M³ and M⁴ are both nitrogen, then p is 2 or 3.


2. The compound of Claim 1 wherein R¹ is selected from:

- 5 (A) aryl;
 (B) substituted aryl, wherein the substituents on said substituted aryl are selected from: (1) halo; or (2) alkyl; or (3) substituted alkyl;
 (C) heteroaryl;
 (D) substituted heteroaryl; or
 10 (E) when R¹ is taken together with X, then the moiety is



3. The compound of Claim 2 wherein R¹ is selected from:

- 15 (A) phenyl;
 (B) substituted phenyl wherein the substituents on said substituted phenyl are selected from: (1) halo; (2) alkyl; (3) alkyl substituted with halo;
 (C) heteroaryl selected from: pyridyl, thienyl, pyrimidinyl, thiazolyl or pyridyl N-Oxide;
 (D) alkyl substituted thiazolyl; or
 20 (E) when R¹ is taken together with X, then the moiety is




wherein c is 0 or 1, and when c is 1 then R⁶ is halo.



4. The compound of Claim 3 wherein R¹ is selected from:

- 25 (A) phenyl;
 (B) substituted phenyl, wherein the substituents on said substituted phenyl are independently selected from: chloro, fluoro or trifluoromethyl;
 (C) heteroaryl selected from:

- 136 -



(D) substituted heteroaryl of the formula:



(E) when R¹ is taken together with X, then the moiety iswherein c is 0 or 1, and when c is 1 then R⁶ is fluoro.5. The compound of Claim 1 wherein R¹ is selected from:

(A) phenyl;

(B) substituted phenyl, wherein the substituents on said substituted phenyl are independently selected from: chloro, fluoro or trifluoromethyl;


(C) pyridyl; or

(D) substituted heteroaryl of the formula:

(E) when R¹ is taken together with X, then the moiety iswherein c is 0 or 1, and when c is 1 then R⁶ is fluoro.6. The compound of Claim 5 wherein R¹ is pyridyl.

- 137 -

7. The compound of Claim 6 wherein R^1 is



8. The compound of Claim 1 wherein X is $=C(NOR^3)$, and R^3 is selected from H or alkyl.

9. The compound of Claim 8 wherein R^3 is selected from H, methyl or ethyl.

10. The compound of Claim 9 wherein R^3 is methyl.

11. The compound of claim 1 wherein: (1) M^2 is nitrogen; and (2) M^3 and M^4 are selected such that: (a) one is carbon and the other is nitrogen, or (b) both are nitrogen.

12. The compound of Claim 11 wherein:

M^3 is carbon, and M^4 is nitrogen;

n is 2;

a is 0 or 1;

b is 0 or 1;

c is 0 or 1, and when c is 1 then R^6 is halo;


e is 1 to 5; and

p is 2.

Y is $=C(O)$; and

Z is C_1 to C_3 alkyl.

13. The compound of Claim 1 wherein Z is



- 138 -


14. The compound of Claim 1 wherein R^2 is a six membered heteroaryl ring.

15. The compound of Claim 14 wherein R^2 is selected from pyridyl, pyridyl substituted with $-NR^4R^5$, pyrimidinyl, or pyrimidinyl substituted with $-NR^4R^5$.

5

16. The compound of Claim 15 wherein R^2 is pyridyl substituted with $-NH_2$, or pyrimidinyl substituted with $-NH_2$.

17. The compound of Claim 16 wherein R^2 is



10

18. The compound of Claim 1 wherein R^4 is H or lower alkyl; R^5 is H, C_1 to C_6 alkyl, or $-C(O)R^4$; R^{12} is alkyl, hydroxy or fluoro; and R^{13} is alkyl, hydroxy or fluoro.

15

19. The compound of Claim 18 wherein R^4 is H or methyl; R^5 is H or methyl; R^{12} is hydroxy or fluoro; and R^{13} is hydroxy or fluoro.

20. The compound of Claim 1 wherein:

(1) R^1 is selected from:

20

(A) aryl;


(B) substituted aryl, wherein the substituents on said substituted aryl are selected from: (1) halo; or (2) alkyl; or (3) substituted alkyl;

(C) heteroaryl; or

(D) substituted heteroaryl; or

25

(E) when R^1 is taken together with X, then the moiety is




(2) X is $=C(NOR^3)$;

- 139 -

- (3) R^3 is selected from H or alkyl;
 (4) M^2 is nitrogen;
 (5) Y is $=C(O)$;
 (6) M^3 and M^4 are selected such that: (1) one is carbon and the other is
 5 nitrogen, or (2) both are nitrogen;
 (7) Z is C_1 to C_3 alkyl; and
 (8) R^2 is a six membered heteroaryl ring.


21. The compound of Claim 16 wherein:
 10 (1) R^1 is selected from:
 (A) phenyl;
 (B) substituted phenyl wherein the substituents on said substituted phenyl are selected from: (1) halo; (2) alkyl; (3) alkyl substituted with halo;
 (C) heteroaryl selected from: pyridyl, thienyl, pyrimidinyl, thiazolyl or
 15 pyridyl N-Oxide; or
 (D) alkyl substituted thiazolyl; or
 (E) when R^1 is taken together with X, then the moiety is




- wherein c is 0 or 1, and when c is 1 then R^6 is halo;
 20 (2) R^3 is selected from H, methyl or ethyl;
 (3) n is 2,
 (4) a is 0 or 1,
 (5) b is 0 or 1,
 (6) c is 0 or 1 and when c is 1 then R^6 is halo,
 25 (7) e is 1 to 5,
 (8) p is 2,
 (9) R^4 is H or lower alkyl,
 (10) R^5 is H, C_1 to C_6 alkyl, or $-C(O)R^4$;
 (11) R^{12} is alkyl, hydroxy or fluoro, and
 30 (12) R^{13} is alkyl, hydroxy or fluoro.

- 140 -

22. The compound of Claim 21 wherein R^2 is



R^1 is




M^2 is nitrogen, M^3 is carbon, and M^4 is nitrogen.

23. The compound of Claim 1 selected from: Compound 23, 30, 31, 32, 33,
 10 41, 44, 45, 49, 50, 52, 53, 54, 55, 56, 57A, 59, 65, 75, 76, 80, 82, 83, 88, 92, 99, 104,
 105, 110, 111, 117, 121, 123, 127, 128, 200-241, 244-273, 275, 278-282, 287, 296,
 301-439, or 446.


24. The compound of Claim 1 selected from: Compound 32, 54, 55, 253A,
 15 287 or 320.

25. The compound of Claim 1 having the formula:




- 141 -

26. The compound of Claim 1 having the formula:




27. The compound of Claim 1 having the formula:




5

28. The compound of Claim 1 having the formula:




10

29. The compound of Claim 1 having the formula:




- 142 -

30. The compound of Claim 1 having the formula:




31. The compound of Claim 1 having the formula:



5

32. The compound of Claim 1 having the formula:



10 33. A pharmaceutical composition comprising: (a) an effective amount of a compound of any of Claims 1 to 32 and a pharmaceutically effective carrier; or (b) an effective amount of a compound of any of Claims 1 to 32, and an effective amount of H_1 receptor antagonist, and a pharmaceutically effective carrier.

15 34. A use of a compound of any of claims 1 to 32 for the manufacture of a medicament for treating: allergy, allergy-induced airway responses, congestion, cardiovascular disease, diseases of the GI tract, hyper and hypo motility and acidic secretion of the gastro-intestinal tract, obesity, sleeping disorders, disturbances of the

- 143 -

central nervous system, attention deficit hyperactivity disorder, hypo and hyperactivity of the central nervous system, Alzheimer's disease, schizophrenia, and migraine.

35. The use of Claim 34 wherein allergy-induced airway responses are
5 treated.

36. The method of Claim 34 wherein allergy or nasal congestion is treated.

37. The use of a compound of any of Claims 1 to 32 and the use of an H₁
10 receptor antagonist for the manufacture of a medicament for treating: allergy, allergy-induced airway responses, and congestion.

38. The use of Claim 37 wherein said H₁ receptor antagonist is selected
from: astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine,
15 chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine,
descarboethoxyloratadine, diphenhydramine, doxylamine, dimethindene, ebastine,
epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine,
meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast,
pyrilamine, promethazine, terfenadine, tripeleminamine, temelastine, trimeprazine or
20 triprolidine.

39. The use of Claim 38 wherein said H₁ receptor antagonist is selected
from: loratadine, descarboethoxyloratadine, fexofenadine or cetirizine.

40. The use of Claim 39 wherein said H₁ receptor antagonist is selected
25 from: loratadine or descarboethoxyloratadine.

30

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 April 2002 (25.04.2002)


PCT

(10) International Publication Number
WO 02/032893 A3

- (51) International Patent Classification⁷: C07D 401/14, 413/14, 401/12, 409/14, 498/04, 417/14, 213/73, A61K 31/4545, A61P 37/08
- (74) Agent: JEANETTE, Henry, C.; Schering-Plough Corporation, Patent Department - K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).
- (21) International Application Number: PCT/US01/32151
- (22) International Filing Date: 15 October 2001 (15.10.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/240,901 17 October 2000 (17.10.2000) US
- (71) Applicant: SCHERING CORPORATION [US/US]; Patent Department - K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).
- (72) Inventors: ASLANIAN, Robert, G.; 144 Philip Drive, Rockaway, NJ 07866 (US). SHIH, Neng-Yang; 1 Maple Drive, North Caldwell, NJ 07006 (US). TING, Pauline, C.; 74 Delwick Lane, New Providence, NJ 07974 (US). BERLIN, Michael, Y.; 14 Hendrick Road, Flemington, NJ 08822 (US). ROSENBLUM, Stuart, B.; 16 Steven Terrace, West Orange, NJ 07052 (US). MC CORMICK, Kevin, D.; 5 Pace Drive, Edison, NJ 08820 (US). TOM, Wing, C.; 133 Cedar Grove Parkway, Cedar Grove, NJ 07009 (US). BOYCE, Christopher, W.; 71 Bonnell Street, Flemington, NJ 08822 (US). MANGIARACINA, Pietro; 4 Montclair Avenue, Monsey, NY 10952 (US). MUTAHI, Mwangi, Wa; 45 Snyder Road, Fords, NJ 08863 (US). PIWINSKI, John, J.; 6 Saddle Ridge Drive, Clinton Township, NJ 08833 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LU, LV, MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UZ, VN, YU, ZA.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- (88) Date of publication of the international search report:
22 August 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

WO 02/032893 A3

(54) Title: PIPERIDINE COMPOUNDS AS ANTI-ALLERGIC



various diseases or conditions, such as, for example, allergy, allergy-induced airway responses, and congestion (e.g., nasal congestion) using the compounds of Formula (I) in combination with a H1 receptor antagonist.

(57) Abstract: Disclosed are novel compounds of the formula: (I) Also disclosed are pharmaceutical compositions comprising the compounds of Formula (I). Also disclosed are methods of treating various diseases or conditions, such as, for example, allergy, allergy-induced airway responses, and congestion (e.g., nasal congestion) using the compounds of Formula (I). Also disclosed are methods of treating

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/32151

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/14 C07D413/14 C07D401/12 C07D409/14 C07D498/04
 C07D417/14 C07D213/73 A61K31/4545 A61P37/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 99 24405 A (ASLANIAN ROBERT G ;MCCORMICK KEVIN D (US); SCHERING CORP (US); MUT) 20 May 1999 (1999-05-20) cited in the application abstract	1, 33, 34, 37
A	WO 97 19074 A (HOECHST MARION ROUSSEL INC) 29 May 1997 (1997-05-29) abstract	1, 33, 34, 37

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the International filing date but later than the priority date claimed

- *T* later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

21 June 2002

Date of mailing of the international search report

01/07/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Diederer, J

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 01/32151

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 1-7 (partially), 11-22 (partially), 33-40 (partially)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-7 (partially), 11-22 (partially), 33-40 (partially)

Present claims 1-7, 11-22 and 33-40 relate to an extremely large number of possible compounds, compositions and uses. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds, compositions and uses claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds of Formula I, in which $X = C(NOR3)$, as defined in claim 8. The search is complete for all examples in the description.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

In ☐ International Application No

PCT/US 01/32151

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9924405	A	20-05-1999	AT 219062 T	15-06-2002
			AU 1296699 A	31-05-1999
			CA 2308586 A1	20-05-1999
			CN 1285822 T	28-02-2001
			EP 1028947 A1	23-08-2000
			HU 0004292 A2	28-11-2001
			JP 2001522835 T	20-11-2001
			TW 421645 B	11-02-2001
			WO 9924405 A1	20-05-1999
			ZA 9810176 A	06-05-1999
WO 9719074	A	29-05-1997	AU 703701 B2	01-04-1999
			AU 1050897 A	11-06-1997
			CA 2237971 A1	29-05-1997
			CN 1202894 A	23-12-1998
			EP 0920425 A1	09-06-1999
			HU 9900927 A2	28-03-2000
			IL 124396 A	19-03-2001
			JP 11513991 T	30-11-1999
			NO 982238 A	01-07-1998
			NZ 323543 A	29-06-1999
			TW 394771 B	21-06-2000
			WO 9719074 A1	29-05-1997
			ZA 9609484 A	10-06-1997
			US 6211199 B1	03-04-2001